Gastrointestinal, Hepatobiliary, and Pancreatic Pathology

β -Caryophyllene Inhibits Dextran Sulfate Sodium-Induced Colitis in Mice through CB2 Receptor Activation and PPAR γ Pathway

Allisson Freire Bento, Rodrigo Marcon, Rafael Cypriano Dutra, Rafaela Franco Claudino, Maíra Cola, Daniela Ferraz Pereira Leite, and João B. Calixto

From the Department of Pharmacology, Biological Sciences Center, Federal University of Santa Catarina Florianópolis, Santa Catarina, Brazil

Cannabinoid receptor 2 (CB2) activation is suggested to trigger the peroxisome proliferator-activated receptor- γ (PPAR γ) pathway, and agonists of both receptors improve colitis. Recently, the plant metabolite (*E*)- β -caryophyllene (BCP) was shown to bind to and activate CB2. In this study, we examined the anti-inflammatory effect of BCP in dextran sulfate sodium (DSS)-induced colitis and analyzed whether this effect was mediated by CB2 and PPAR γ . Oral treatment with BCP reduced disease activity, colonic macro- and microscopic damage, myeloperoxidase and *N*-acetylglucosaminidase activities, and levels and mRNA expression of colonic tumor necrosis factor- α , IL-1 β , interferon- γ , and keratinocyte-derived chemokine. BCP treatment also inhibited the activation of extracellular signal-regulated kinase 1/2, nuclear factor κ B, I κ B-kinase α/β , cAMP response element binding and the expression of caspase-3 and Ki-67. Moreover, BCP enhanced IL-4 levels and forkhead box P3 mRNA expression in the mouse colon and reduced cytokine levels (tumor necrosis factor- α , keratinocyte-derived chemokine, and macrophage-inflammatory protein-2) in a culture of macrophages stimulated with lipopolysaccharide. The use of the CB2 antagonist AM630 or the PPAR γ antagonist GW9662 significantly reversed the protective effect of BCP. Confirming our results, AM630 reversed the beneficial effect of BCP on pro-inflammatory cytokine expression in IEC-6 cells. These results demonstrate that the anti-inflammatory effect of BCP involves CB2 and the PPAR γ pathway and suggest BCP as a possible therapy for the treatment of inflammatory bowel disease. (*Am J Pathol* 2011, 178:1153–1166; DOI: 10.1016/j.ajpath.2010.11.052)

Inflammatory bowel diseases (IBDs) are a group of chronic diseases that affect the gastrointestinal tract and have been mainly subdivided as ulcerative colitis and Crohn's disease.¹ The IBDs are characterized by a strong leukocyte activation and infiltration into the intestinal tissues, the release of proinflammatory cytokines² and enzymes, and the formation of reactive oxygen species. All of these events can induce an extensive and unbalanced activation of the mucosal immune system, driven by the commensal flora.³

Recent evidence suggests a role for the cannabinoid system in IBD regulation. Cannabinoid receptors 1 and 2 (CB1 and CB2) are expressed in normal human colon^{4,5} and are up-regulated in IBD colonic tissue. In addition, an enhanced level of endocannabinoids was found in biopsy specimens from patients with ulcerative colitis.⁵ It is thought that CB1 activation results in a decrease of intestinal hypermotility and hypersecretion, whereas the activation of CB2 results in the inhibition of proinflammatory mediators. In addition, both CB2 and CB1 knockout mice are more susceptible to the development of experimental colitis, and the activation of these receptors is extremely important for the abrogation of intestinal inflammation.⁶ The role of CB2, however, is directly involved with innate immune system, because CB2 is primarily expressed in immune cells, such as macrophages, CD4⁺ and CD8⁺ T cells, monocytes, and polymorphonuclear neutro-

Supported by the National Council for Scientific and Technological Development (CNPq: Conselho Nacional de Desenvolvimento Científico e Tecnológico) (to A.F.B., R.M. and R.C.D., R.F.C., M.C. and D.F.P.L), Coordination for Improvement of Higher Education (CAPES: Coordenação de Aperfeiçoamento de Pessoal de Nível Superior), and the Foundation for Research Support-State of Santa Catarina (FAPESC: Fundação de Apoio à Pesquisa Científica Tecnológica do Estado de Santa Catarina) to A.F.B., R.M., R.C.D., R.F.C., and M.C. (who received grants from CNPq).

A.F.B. and R.M. contributed equally to this study.

Accepted for publication November 23, 2010.

Supplemental material for this article can be found at <http://ajp.amjpathol.org> or at doi:10.1016/j.ajpath.2010.11.052.

Address reprint requests to João B. Calixto, Ph.D., Departamento de Farmacologia, Centro de Ciências Biológicas, Universidade Federal de Santa Catarina (UFSC), Campus Universitário, Trindade, 88049-900, Florianópolis, SC, Brazil. E-mail: calixto@farmaco.ufsc.br or calixto3@terra.com.br.

phils,^{7,8} which are intimately related to the inflammatory response in IBD.^{9,10} Although CB2 expression has been recently reported within the central nervous system,¹¹ including the spinal cord, microglial cultures, brainstem, and cortex,¹² it has not been associated with the central nervous system side effects of *CB1* activation. This observation suggests that CB2 is a more attractive therapeutic target for the treatment of IBDs, such as ulcerative colitis.

Several studies support the notion that endocannabinoids exert their anti-inflammatory properties, at least in part, by the activation of the peroxisome proliferator-activated receptor-gamma (PPAR γ) pathway.^{13,14} PPAR γ is a member of the superfamily of nuclear receptors and has important anti-inflammatory activity,^{15,16} because it inhibits the activation of nuclear factor κ B (NF κ B) and the expression of the proinflammatory cytokines IL-1 β and tumor necrosis factor- α (TNF- α).^{17–19} The colon expresses a high density of PPAR γ ,^{20,21} and much evidence points to the involvement of the PPAR γ receptor in the regulation of intestinal inflammation.^{20,22}

Recently, Gertsch et al²³ demonstrated that the sesquiterpene (*E*)- β -caryophyllene (BCP) selectively binds to CB2 and acts as a full agonist. In confirming this hypothesis, the authors demonstrated that oral administration of this compound exerted potent anti-inflammatory effects in wild-type mice but not in CB2 knockout (*Cnr2*^{-/-}) mice. However, the mechanisms underlying the anti-inflammatory effects of BCP are not completely understood. BCP could inhibit the pathways triggered by the activation of the Toll-like receptor complex CD14/TLR4/MD2, which typically leads to the expression of proinflammatory cytokines (IL-1 β , IL-6, IL-8, and TNF- α).²⁴ In addition, BCP suppresses extracellular signal-regulated kinase 1/2 (ERK1/2) and c-Jun N-terminal kinase 1/2 (JNK1/2) signaling. All of these effects contribute to the anti-inflammatory effects of the sesquiterpene.²³ BCP is present in a number of plant species that have been widely used in folk medicine for their anti-inflammatory and healing properties, including wild sage [*Cordia curassavica* (Jacq.) Roem. & Schult. (synonym: *Cordia verbenacea* DC.)] and black pepper (*Piper nigrum* L.).^{25,26}

Because both PPAR γ and the cannabinoid system are relevant targets for the treatment of IBD, we investigated the molecular mechanisms underlying the anti-inflammatory effects of oral BCP in experimental colitis induced by dextran sulfate sodium (DSS) administration in mice. The findings suggest that BCP seems to reduce intestinal inflammation through CB2 activation and acts directly or indirectly with the PPAR γ nuclear receptor. We therefore suggest BCP as a possible therapy for the treatment of IBD.

Materials and Methods

Animals

Male CD1 mice (8 to 10 weeks of age) were obtained from the animal housing unit of the Laboratory of Experimental Pharmacology at the Federal University of Santa Catarina (Laboratório de Farmacologia Experimental–LAFEX, Universidade Federal de Santa Catarina Florianópolis, SC, Brazil). Animals were housed in collective

cages at 22 $\pm$ 1°C under a 12-hour light/dark cycle (lights on at 7:00 a.m.) with free access to laboratory chow and drinking water (tap water). Experiments were performed during the light phase of the cycle. The experimental procedures were previously approved by the Federal University of Santa Catarina Committee on the Ethical Use of Animals and the study was conducted in accordance with Brazilian regulations on animal welfare.

Induction and Assessment of DSS-Induced Colitis

Male CD1 mice (5 to 7 per group) were provided with a solution of filtered water containing 3% (w/v) DSS (mol. wt. 36,000 to 50,000 kDa) *ad libitum* over a 5-day period. Every other day, the DSS solution was replenished and the amount consumed was measured. At the end of this 5-day period, DSS was replaced by normal drinking water for 2 days and, at the end of day 7, the animals were euthanized. Control mice received only normal drinking water. All animals were examined once a day and the disease activity index (DAI) was assessed as described previously.^{27,28} Briefly, the DAI combines scores for weight loss, stool consistency, and bleeding. Scores were defined as follows: stool consistency was graded 0 for no diarrhea, 2 for loose stool that did not stick to the anus, and 4 for liquid stool that did stick to the anus. The presence of fecal blood was graded 0 for none, 2 for moderate, and 4 for gross bleeding. For weight loss, a value of 0 was assigned if body weight increased or remained within 1% of baseline, 1 for a 1% to 5% loss, 2 for a 5% to 10% loss, 3 for a 10% to 15% loss, and 4 for weight loss >15%. At the end of the 7-day period, the colons were removed and examined for weight, the consistency of the stool found within, and gross macroscopic appearance and length (which was measured from 1 cm above the anus to the top of the cecum). This macroscopic scoring was performed as described previously²⁹ (for detail, see Supplemental Table S1 at <http://ajp.amjpathol.org>). For each animal, the three subscores were summed to provide the total macroscopic score.

In another set of experiments, each excised portion of the distal colon was fixed immediately in a 4% (w/v) formaldehyde solution. Tissues were embedded in paraffin, sectioned (5 μ m thickness), mounted on glass slides, and deparaffinized. For a general histological analysis, slices were stained using standard H&E techniques. Samples were analyzed by light microscopy and were scored as described previously²⁹ (see Supplemental Table 2 at <http://ajp.amjpathol.org>). In this case, the experiments were performed in a double-blind manner.

Induction and Assessment of Oxazolone-Induced Colitis

CD1 mice were presensitized by the epicutaneous application of oxazolone [4-ethoxymethylene-2-phenyl-2-oxazolin-5-one, 3% (w/v) in 150 μ L 100% ethanol] on day 7. Briefly, 1-day fasted mice were given oxazolone [3% in 100 μ L 50% (v/v) ethanol] delivered using a polyethylene PE50 catheter that was inserted into the colon 4 cm

proximal to the anus under anesthesia administered with an intraperitoneal injection of a mixture of ketamine and xylazine. Control mice received 100 μ L of 50% ethanol under the same conditions. After colonic instillation, the animals were kept in a head-down position for 2 minutes and were kept from food and water for 4 hours (in a separate holding cage) before transfer to their home cages. Body weight was estimated daily for 72 hours after treatment, starting on the day preceding colitis induction. The number of surviving animals in each group was recorded daily to determine survival rates.

Pharmacological Treatments

Animals were orally treated by gavage with 12.5, 25, or 50 mg/kg of BCP twice a day from day 0 to day 7 (preventive treatment) or with 50 mg/kg from day 3 to day 7 (therapeutic treatment). In another set of experiments, animals received an intraperitoneal injection of the CB2 antagonist AM630 (10 mg/kg) or the PPAR γ antagonist GW9662 (1 mg/kg) once a day for 7 days, given 30 minutes before the BCP (50 mg/kg). The dose of each drug was chosen based on preliminary studies or previous publications.^{30–32} The BCP was solubilized in a 0.9% NaCl solution and the AM630 and GW9662 were prepared in 2% dimethyl sulfoxide in saline solution. All-vehicle solutions were used for the respective control animal treatments.

To evaluate the potential effects of BCP on oxazolone-induced colitis, mice were treated twice daily for 3 days with either BCP (50 mg/kg per day, p.o.) or vehicle, starting 24 hours after oxazolone administration. All animals were sacrificed by cervical dislocation at 72 hours after oxazolone administration (ie, 4 hours after receiving the last treatment).

Myeloperoxidase and N-Acetylglucosaminidase Assays

Neutrophil and macrophage infiltration into the colon was assessed indirectly by measuring myeloperoxidase (MPO) and *N*-acetylglucosaminidase (NAG) activities, respectively. The MPO and NAG assays were performed as described previously.⁹ On day 7, animals were sacrificed and colon tissue segments were homogenized in 5% EDTA/NaCl buffer (pH 4.7) and centrifuged at 10,000 $\times g$ for 15 minutes at 4°C. The pellet was resuspended in 0.5% hexadecyl trimethyl ammonium bromide buffer (pH 5.4), and the samples were frozen in liquid nitrogen and thawed three times. On final thawing, the samples were similarly centrifuged, and 25 μ L of the supernatant was used for MPO and NAG assays. The MPO enzymatic reaction was assessed by the addition of 1.6 mmol/L tetramethylbenzidine, 80 mmol/L NaPO $_4$ and 0.3 mmol/L hydrogen peroxide (H $_2$ O $_2$). The NAG activity measurement was determined by adding 2.25 mmol/L *p*-nitrophenyl-2-acetamido- β -glucopyranoside and 100 μ L 50 mmol/L citrate buffer (pH 4.5). The absorbances were measured spectrophotometrically at 690 and 405 nm for MPO and NAG, respectively, and the results are expressed in optical density per milligram tissue.

Determination of Cytokine Levels

For determination of cytokine levels, colon segments were homogenized in phosphate buffer containing 0.05% Tween 20, 0.1 mmol/L phenylmethylsulfonyl fluoride, 0.1 mmol/L benzethonium chloride, 10 mmol/L EDTA, and 20 UI aprotinin A. The homogenate was centrifuged at 3000 $\times g$ for 10 minutes, and the supernatants were stored at -70°C for further analysis. Levels of TNF- α , IL-1 β , keratinocyte-derived chemokine (CXCL1/KC), interferon- γ (INF- γ), IL-4, IL-10, and transforming growth factor- β were evaluated using enzyme-linked immunosorbent assay kits from R&D Systems (Minneapolis, MN) according to the manufacturer's instructions. The amount of protein in each sample was measured using the Bradford method.³³

RNA Extraction and Real-Time PCR

Total RNA from colons was extracted using the TRIzol protocol, and its concentration was determined using a NanoDrop 1100 spectrophotometer (NanoDrop Technologies, Wilmington, DE). A reverse transcription assay was performed as described in the M-MLV reverse transcriptase protocol according to the manufacturer's instructions (Applied Biosystems, Foster City, CA). cDNA (300 ng) was amplified in triplicate using the TaqMan universal PCR master mix kit with specific TaqMan gene expression target genes, the 3' quencher MGB and FAM-labeled probes for mouse TNF- α (Mm00443258_m1), IL-1 β (Mm01336189_m1), CXCL1/KC (Mm00433859_m1), IFN- γ (Mm99999071_m1), CB2 (Mm00438286_m1), PPAR γ (Mm00440940_m1), forkhead box P3 (Foxp3) (Mm00475165_m1), GAPDH (NM_008084.2), rat TNF- α (Rn99999017_m1), and rat β actin (Rn00667869_m1) that was used as an endogenous control for normalization. The PCR reactions were performed in a 96-well optical reaction plate (Applied Biosystems, Foster City, CA). The thermocycler parameters were as follows: 50°C for 2 minutes, 95°C for 10 minutes, 50 cycles of 95°C for 15 seconds, and 60°C for 1 minute. Expression of the target genes was calibrated against conditions found in control animals (ie, animals that received vehicle).

Mesenteric Lymph Node Macrophage Purification and Stimulation

Mesenteric lymph nodes obtained from naïve mice were macerated in RPMI 1640 medium and filtered through a 220- μ m filter. The resulting suspension was centrifuged at 1500 $\times g$ for 7 minutes, the supernatant was discarded, and the cell pellet was resuspended in RPMI 1640 medium supplemented with 10% fetal bovine serum, 20 mmol/L HEPES, 3 $\times 10^{-5}$ mol/L 2-mercaptoethanol, 100 U/ml penicillin, and 100 μ g/ml streptomycin. Macrophages were purified by adherence to plastic. Briefly, 0.1 ml of the cell suspension (1 $\times 10^6$ cells/ml) were distributed in 96-well plates and incubated for 2 hours at 37°C in a 5% CO $_2$ enriched atmosphere. Afterward, the cells were washed and the remaining cells were stimulated for 24 hours with lipopolysaccharide (LPS; 10 μ g/ml) in the presence or absence of BCP (10 μ mol/L) in a final volume of 200 μ L/well.

After stimulation, the plate was centrifuged ($200 \times g$, 10 minutes), and the cell-free supernatant was collected and stored at -70°C for cytokine determination. The concentration of TNF- α , IL-4, MIP-2, and CXCL1/KC was measured using an enzyme-linked immunosorbent assay kit according to the manufacturer's instructions.

Murine Bone Marrow-Derived Macrophages

CD1 mice were sacrificed by cervical dislocation. Total bone marrow was obtained from mice by flushing the femurs and tibiae with Dulbecco's modified Eagle's medium. Bone marrow-derived macrophages were obtained as described previously.³⁴ Briefly, bone marrow mononuclear phagocyte precursor cells were propagated in suspension by culturing in macrophage medium (Dulbecco's modified Eagle's medium containing glucose, supplemented with 2 mmol/L L-glutamine, 10% fetal calf serum, 10 mmol/L HEPES, 100 $\mu\text{g}/\text{ml}$ streptomycin, and 100 U/ml penicillin (all from Sigma-Aldrich, St. Louis, MO) supplemented with 20% L929 cell-conditioned medium (as a source of macrophage colony stimulating factor). Cells were incubated at 37°C in a 5% CO_2 enriched atmosphere and were fed on day 5 by replacing the medium supplemented with 20% L929 cell-conditioned medium. Cells were harvested on day 7 and 2×10^5 cells/ml were cultured in a 96-well cell culture plate for 24 hours. Afterward, adherent cells were stimulated for 24 hours with LPS (10 $\mu\text{g}/\text{ml}$) in the presence or absence of BCP (10 $\mu\text{mol}/\text{L}$) in a final volume of 250 $\mu\text{L}/\text{well}$. After stimulation, the plate was centrifuged ($200 \times g$, 10 minutes) and the cell-free supernatant was collected and stored at -70°C for cytokine determination. The concentration of TNF- α and CXCL1/KC was measured using an enzyme-linked immunosorbent assay kit according to the manufacturer's instructions.

Intestinal Epithelial Cell-6 Culture

Intestinal epithelial cell-6 (IEC-6), a nontransformed rat intestinal epithelial-derived cell line, was maintained in Dulbecco's modified Eagle's medium (containing glucose and supplemented with 2 mmol/L L-glutamine, 10 mmol/L HEPES, 100 $\mu\text{g}/\text{ml}$ streptomycin, and 100 U/ml penicillin). Cells (2×10^5 cells/well) were distributed in 24-well plates and all experiments were performed after cultures reached confluence (2 to 3 days after plating). After confluence was achieved, the cell culture medium was replaced. Cells were treated with BCP (10 $\mu\text{mol}/\text{L}$) for 30 minutes before stimulation.

In another set of experiments, to assess the participation of the CB2 in the effect of BCP, cells were pretreated with the CB2 selective antagonist AM630 at 1 hour before BCP treatment. Cells were then stimulated with LPS (1 $\mu\text{g}/\text{ml}$) for 4 hours in the presence or absence of BCP, AM630 (5 $\mu\text{mol}/\text{L}$), or vehicle. Control cells were incubated with the appropriate corresponding vehicles. After stimulation, the plate was centrifuged ($200 \times g$, 10 minutes), and the cell-free supernatant was collected and stored at -70°C for determination of the protein levels for the cytokine CINC-1 and mRNA expression for TNF- α , as described above.

Immunohistochemistry Analysis

Immunohistochemical reaction analysis was performed using the colon according to methods described previously.³⁵ Slices (5 μm thickness) were stained using the following primary antibodies and respective dilutions: polyclonal goat anti-phospho-ERK (1:300), monoclonal mouse anti-phospho-p65 NF κB (1:50), polyclonal rabbit anti-phospho-CREB (1:300), polyclonal rabbit anti-phospho-IKK α/β (1:100), polyclonal rabbit anti-cleaved-caspase-3 (1:500), polyclonal rabbit anti-claudin-4 (1:200), and polyclonal rabbit anti-Ki-67 (1:300). High-temperature antigen retrieval was performed by the immersion of the slides in a water bath at 95°C to 98°C in a 10 mmol/L trisodium citrate buffer of pH 6.0 for 45 minutes. The nonspecific binding was blocked by incubating sections for 1 hour with goat normal serum diluted in PBS. After an overnight incubation at 4°C with primary antibodies, the slides were washed with PBS and incubated with the secondary antibody Envision Plus (ready-to-use) for 1 hour at room temperature. After incubation with the appropriate biotinylated secondary antibody, the sections were developed with 3,3'-diaminobenzidine (DakoCytomation, Glostrup, Denmark) in a chromogen solution for an exact amount of time and counterstained with Harris hematoxylin.

Images of colon sections stained with antibodies to phospho-ERK, phospho-p65 NF κB , phospho-CREB, phospho-IKK α/β , cleaved caspase-3, claudin-4, and Ki-67 were acquired using a Sight DS-5M-L1 digital camera connected to an Eclipse 50i light microscope (both from Nikon, Melville, NY) and image acquisition software (Qcapture Pro 5.1; QImaging, Surrey, BC, Canada). All image analysis was performed using NIH ImageJ 1.36b imaging software (NIH, Bethesda, MD). Total pixel intensity was determined for phospho-ERK, phospho-p65 NF κB , phospho-CREB, phospho-IKK α/β , claudin-4, and Ki-67, and data were reported as optical density. To analyze the apoptosis index in the colonic tissue after DSS-administration, cleaved caspase-3 positive cells were assessed by visual inspection by counting the labeled cells in a colon section using a counting grid at $\times 400$ magnification.

Drugs and Reagents

Dextran sulfate sodium (DSS) (mol. wt. 36,000 to 50,000 kDa) was obtained from MP Biomedicals (Solon, OH). Both BCP and oxazolone were obtained from Sigma-Aldrich. GW9662 was obtained from Calbiochem (Merck, Darmstadt, Germany). AM630 was obtained from Tocris Bioscience (Ellisville, MO). Formaldehyde was obtained from Merck (Darmstadt, Germany). Polyclonal goat anti-phospho-ERK (pERK), monoclonal mouse anti-phospho-p65 (NF κB), polyclonal rabbit anti-nuclear phospho-cyclic AMP response element binding protein (CREB), polyclonal rabbit anti-IKK α/β , and polyclonal rabbit anti-cleaved-caspase-3 were purchased from Cell Signaling Technology (Danvers, MA). Polyclonal rabbit anti-claudin-4 was purchased from Spring Bioscience (Pleasanton, CA). Polyclonal rabbit anti-Ki-67 was purchased from Abcam (Cambridge, MA). Secondary anti-

body Envision Plus, streptavidin–horseradish peroxidase reagent and 3,3'-diaminobenzidine chromogen were purchased from DakoCytomation (Carpinteria, CA). Hydrogen peroxide, Tween 20, Tween 80, EDTA, aprotinin, phosphate-buffered saline, eosin, hematoxylin, tetramethylbenzidine, hydrogen peroxide, and *o*-phenylenediamine dihydrochloride were purchased from Sigma-Aldrich. Mouse CXCL1/KC, MIP-2, IFN- γ , TNF- α , IL-4, transforming growth factor- β , IL-10, IL-1 β /IL-1F2, and rat CINC-1 DuoSet kits were obtained from R&D Systems (Minneapolis, MN). TRIzol and M-MLV reverse transcriptase were purchased from Invitrogen (Carlsbad, CA). Primers and probes for mouse TNF- α (Mm00443258_m1), CXCL1/KC (Mm00433859_m1), IFN- γ (Mm99999071_m1), PPAR γ (Mm00440940_m1), FoxP3 (Mm00475165_m1), CB2 (Mm00438286_m1), GAPDH (NM_008084.2), rat TNF- α (Rn99999017_m1), and rat β actin (Rn00667869_m1) TaqMan universal PCR master mix kit were purchased from Applied Biosystems (Foster City, CA).

Data Analysis

All data are expressed as means $\pm$ SEM. For nonparametric data, a Kruskal–Wallis test followed by a Dunn test was used. For parametric data, the statistical differences between groups were determined by one-way analysis of variance followed by a Student–Newman–Keuls test. Statistical analyses were performed using GraphPad Prism 4 software (GraphPad Software, San Diego, CA). $P < 0.05$ was considered statistically significant.

Results

β -Caryophyllene Ameliorates DSS- and Oxazolone-Induced Colitis

Mice treated with 3% DSS developed a severe illness characterized by bloody diarrhea and sustained weight loss. Preventive treatment with BCP in three different doses (12.5, 25, or 50 mg/kg, p.o.) was used to assess the potential dose-dependent effects of BCP. At the two higher doses, 25 and 50 mg/kg, BCP resulted in a significant reduction of the DAI score. The highest dose (50 mg/kg, p.o.) was more effective and exhibited an earlier effect (significant from day 4; Figure 1, A and B). For this reason, a dose of 50 mg/kg (p.o.) of BCP was used in subsequent experiments. Mice treated with BCP (50 mg/kg) were protected from marked weight loss and recovered a healthy appearance, similar to that of control mice (Figure 1C). DSS administration resulted in colon inflammation associated with hyperemia, ulceration, and bowel wall thickening leading to an increase of macroscopic colon damage and colon length reduction (Figure 1, D and E). BCP oral treatment markedly reduced macroscopic damage in a dose-dependent manner (Figure 1D) and partially prevented colon length reduction (Figure 1, E and F) on day 7 after colitis induction. Therapeutic treatment with BCP (50 mg/kg, p.o.) also promoted an improvement in clinical parameters analyzed from day 3. Of note, no difference was observed in water or DSS solution consumption between the

Figure 1. β -Caryophyllene (BCP) ameliorates DSS-induced colitis. Mice received DSS (3%) for 5 days and drinking water for the next 2 days. Animals were orally treated by gavage with 12.5, 25, or 50 mg/kg of BCP twice a day from day 0 to day 7 (preventive treatment) or with 50 mg/kg from day 3 to day 7 (therapeutic treatment). Preventive or therapeutic oral treatment with BCP improved the disease activity index (DAI) score (A and B), reduced body weight loss (C) and colon macroscopic damage (D), and enhanced colon length (E), compared with mice from the DSS group. Data are reported as means $\pm$ SEM ($n = 5$ to 7 mice per group). * $P < 0.05$ versus the DSS-treated group; ** $P < 0.05$ versus the control group; *** $P < 0.05$ versus the preventive BCP-treated group (50 mg/kg, p.o.). (F) Representative photographs of colons from control mice (CTL), DSS-treated mice, and preventive BCP-treated mice (50 mg/kg, p.o.).

experimental groups (data not shown). No side effects were observed with BCP alone (50 mg/kg) in acute toxicological studies; the administration of BCP alone did not alter the DAI score, microscopic damage, or MPO activity in animal colons, compared with the control group (see Supplemental Figure S1 at <http://ajp.amjpathol.org>).

To investigate if BCP treatment would also prove to be beneficial in another preclinical model of colitis, we tested its effects on some parameters of colitis induced by the hapten reagent oxazolone. The rectal administration of oxazolone dissolved in ethanol induced a severe colitis in CD1 mice that was characterized by weight reduction and diarrhea, which led to high death rates. Preventive treatment with BCP (50 mg/kg, p.o., twice daily) markedly enhanced survival and body weight gain (see Supplemental Figure S2 at <http://ajp.amjpathol.org>).

β -Caryophyllene Treatment Inhibits Cellular Influx and Decreases Colon Damage

DSS-induced colon damage is associated with an influx of inflammatory cells, such as neutrophils and macrophages, into the intestinal mucosa.³⁶ We therefore assessed whether the protective effect of BCP treatment in DSS-mediated colitis was associated with alterations in the composition of inflammatory cell populations in the intestinal mucosa. At 7 days after the initiation of DSS treatment, mucosal neutrophil and macrophage infiltration into the colon were indirectly assessed by measuring MPO and NAG activities, respectively. DSS-treated mice displayed a relevant in-

crease in colonic MPO levels and a slight increase in NAG activity, compared with control animals (Figure 2, A and B). Preventive treatment with BCP (12.5, 25, or 50 mg/kg, p.o.) significantly prevented both MPO and NAG activities incrementally (Figure 2, A and B). Similar results were observed with the therapeutic treatment of BCP (50 mg/kg, p.o.).

To further confirm the MPO and NAG analysis, colons were processed for histological observation. Representative sections with H&E staining revealed a marked cell infiltration into the lamina propria and colonic mucosa in colons from DSS-treated mice. In colons from the control group, no histopathological changes were observed. The colonic tissue from DSS-treated mice appeared thick and sometimes ulcerated, showing a distortion of crypts (Figure 2, C and D). Of note, the histological evaluation of colons from BCP-treated mice revealed a pronounced reduction in the inflammatory response, resulting in a decreased microscopic damage score, compared with colons from DSS-treated mice (Figure 2, C and D).

β -Caryophyllene Treatment Reduces the Production and Expression of Inflammatory Mediators in DSS-Induced Colitis

Accumulating data in the literature suggest that cytokines and chemokines (eg, TNF- α , IL-1 β , INF- γ , and CXCL1/KC) are critically involved in inflammatory cell recruitment.^{9,37} We therefore assessed the mRNA expression and the colonic protein levels of these inflammatory mediators on day 7 after induction of colitis. DSS administra-

Figure 2. BCP decreases cell influx and reduces microscopic colon damage. At 7 days after BCP oral treatment, colon tissues were processed for histological evaluation or for the measurement of myeloperoxidase (MPO) activity and *N*-acetylglucosaminidase (NAG) activity. Preventive (12.5, 25, and 50 mg/kg, p.o.) or therapeutic (50 mg/kg, p.o.) treatment with BCP reduced MPO (A) and NAG (B) activity. C: Preventive treatment with BCP (50 mg/kg, p.o.) decreased the microscopic damage score in mouse colon. Data are reported as means $\pm$ SEM ($n = 5$ to 7 mice per group). * $P < 0.05$ versus the DSS-treated group; ** $P < 0.05$ versus the control group. D: Representative paraffin sections of colons from control mice, DSS-treated, and BCP-treated mice (50 mg/kg, p.o.), stained with H&E. Original magnification, $\times 200$.

Figure 3. Treatment with BCP changes colonic protein levels and mRNA expression of inflammatory mediators. At the end of 7 days, colon tissue was collected and processed for cytokine levels and mRNA expression. **A–E:** Enzyme-linked immunosorbent assay. Preventive treatment with BCP (50 mg/kg, p.o.) reduced colonic levels of tumor necrosis factor- α (TNF- α) (**A**), IL-1 β (**B**), keratinocyte-derived chemokine (CXCL1/KC) (**C**), and interferon- γ (INF- γ) (**D**) and restored levels of IL-4 (**E**). **F–J:** Real-time PCR. The same scheme of treatment with BCP also impaired the increase colonic mRNA expression of TNF- α (**F**), IL-1 β (**G**), CXCL1/KC (**H**), and INF- γ (**I**). In addition, BCP treatment increased colonic forkhead box P3 (Foxp3) mRNA expression (**J**) as assessed by real-time PCR. The real-time PCR assay was performed in duplicate and *GAPDH* mRNA was used to normalize the relative amount of mRNA. Data are reported as means $\pm$ SEM ($n = 5$ to 7 mice per group). * $P < 0.05$ versus the DSS-treated group; ** $P < 0.05$ versus the control group.

tion resulted in a pronounced increase in colonic TNF- α , IL-1 β , CXCL1/KC, and INF- γ protein levels (Figure 3, A–D) and mRNA expression (Figure 3, F–I). Preventive treatment with BCP (50 mg/kg) resulted in a striking decrease in these mediators (Figure 3).

The modulation of colonic anti-inflammatory cytokine levels by BCP was also investigated. BCP treatment restored colonic IL-4 levels (Figure 3E), which were reduced in DSS-treated mice, but failed to change IL-10 and transforming growth factor- β levels, (see Supplemental Figure S3 at <http://ajp.amjpathol.org>). Additionally, preventive BCP treatment significantly increased mRNA for the transcription factor FoxP3, compared with the DSS group (Figure 3J). This result suggests that the anti-inflammatory effect displayed by BCP might be related, at least in part, to an increase in CD4⁺ FoxP3⁺ regulatory T cells. Additional experiments are needed to confirm this hypothesis.

β -Caryophyllene Reduces Cytokine Release from LPS-Stimulated Macrophages

Because the inflammatory cells present in the DSS-treated colon produce cytokines and chemokines, the observed reduction of these colonic mediators could simply be a consequence of decreased cell migration.

To clarify whether BCP could also diminish the production of inflammatory mediators or modify the profile of the cytokines produced by the inflammatory cells present in colonic tissue after inflammatory stimulus, we performed a mesenteric lymph node macrophage culture and evaluated the influence of BCP on production of CXCL1/KC, macrophage inflammatory protein-2 (MIP-2), TNF- α , and IL-4 after *in vitro* stimulation with LPS (10 μ g/ml, for 24 hours), which is an important component of colitis-induced damage. LPS stimulation markedly increased levels of CXCL1/KC, MIP-2, and TNF- α (Figure 4, A–C). This *in vitro* treatment with BCP (10 μ mol/L) significantly reduced levels of CXCL1/KC, MIP-2, and TNF- α after administration of LPS (Figure 4, A–C). However, BCP (10 μ mol/L) failed to increase IL-4 levels in the macrophage culture after LPS administration (see Supplemental Figure S4 at <http://ajp.amjpathol.org>).

To investigate whether BCP treatment (10 μ mol/L) could also reduce proinflammatory cytokine levels in another macrophage culture, we evaluated its effects on murine bone marrow-derived macrophages. Production of CXCL1/KC and TNF- α by LPS-stimulated bone marrow-derived macrophages was significantly reduced by pretreatment (30 minutes) with BCP (Figure 4, D and E).

Figure 4. BCP reduces proinflammatory cytokine production in macrophages stimulated with LPS. Macrophages from mesenteric lymph nodes (**A–C**) and bone marrow (**D** and **E**) of naïve mice were stimulated with LPS (10 μ g/well) in the presence or absence of BCP (10 μ mol/L per well) for 24 hours, and the culture supernatants were analyzed for cytokine levels. BCP incubation reduced production of CXCL1/KC (**A** and **D**), macrophage inflammatory protein-2 (MIP-2) (**B**), and TNF- α (**C** and **E**). Data are reported as means $\pm$ SEM ($n = 5$ /group). * $P < 0.05$ versus the LPS-treated group; ** $P < 0.05$ versus the control group.

β -Caryophyllene Inhibits Colonic NF κ B, CREB, ERK1/2, and IKK α/β Activation Induced by DSS

Mitogen-activated protein kinases (MAPK) and transcriptional factors (eg, NF κ B and CREB) are among the major proinflammatory signaling pathways involved in colitis.^{35,38} We therefore further investigated whether BCP treatment could prevent the activation of the p65 NF κ B subunit, CREB, ERK1/2, and IKK α/β . No significant activation of NF κ B, CREB, ERK1/2, or IKK α/β was observed in colons from saline-treated animals (control group). Nonetheless, pronounced phosphorylation of the p65 NF κ B subunit, CREB, ERK1/2, and IKK α/β were detected 7 days after beginning the treatment with DSS (Figure 5, A–D; see also Supplemental Figure S5 at <http://ajp.amjpathol.org>). Of note, oral treatment with BCP consistently impaired activation of this signaling pathway.

β -Caryophyllene Inhibits Colonic Caspase-3 and Ki-67 Activation Induced by DSS but Not Claudin-4

Caspase-3, Ki-67, and claudin-4 play a central role in the inflammatory process. Caspase-3 is important to cell apoptosis.³⁹ Ki-67 is a nuclear protein necessary for cell proliferation,⁴⁰ and claudin-4 is a fundamental component of tight junctions and adhesion cells.⁴¹ To analyze the role of BCP in apoptosis and in cell proliferation and adhesion, we evaluated cleaved caspase-3, Ki-67, and claudin-4 staining in colon tissue after DSS administration. Tissue sections from control mice exhibited very low levels of specific staining for cleaved caspase-3 and Ki-67 in epithelial cells (Figure 5, E and F). In contrast, at 7 days after DSS administration, the DSS group showed intense immunostaining for cleaved caspase-3 and Ki-67

expression in colon tissue; however, treatment with BCP (50 mg/kg, p.o., b.i.d.) during the 7 days after DSS administration significantly inhibited apoptosis and the proliferation index in colon tissue, as measured by monoclonal antibodies against cleaved caspase-3 and Ki-67, respectively (Figure 5, E and F; see also Supplemental Figure S6 at <http://ajp.amjpathol.org>). In addition, DSS-induced colitis was associated with significant decreases in the colonic expression of the tight junction protein claudin-4, but treatment with BCP did not restore the levels of this protein (see Supplemental Figure S7 at <http://ajp.amjpathol.org>) or immunostaining (see Supplemental Figure S6 at <http://ajp.amjpathol.org>).

Beneficial Effects of β -Caryophyllene on DSS-Induced Colitis Are Reversed by CB2 and PPAR γ Selective Antagonists

Previous studies have demonstrated that CB2 is up-regulated in inflamed colons of patients with colitis.^{29,42} The present results corroborate these data by showing a great increase of CB2 mRNA expression in colonic tissue from DSS-treated mice, compared with colons from the control group (Figure 6E). Of note, preventive oral treatment with BCP markedly diminished CB2 expression in colons from mice on day 7 after DSS treatment (Figure 6E).

BCP has been shown previously to be a selective agonist of cannabinoid receptor 2.²³ Importantly, CB2 activation decreases colon inflammation in different experimental models.^{29,43} We investigated whether a CB2 selective antagonist could prevent the anti-inflammatory effect of BCP. For this purpose, mice were treated once a day with the selective CB2 antagonist AM630 (10 mg/kg, p.o.), alone or in combination with BCP, for 7 days. Treatment with AM630 alone did not alter the inflammatory parameters observed in the DSS-treated group (Figure 6,

Figure 5. BCP treatment impairs activation of NF κ B, CREB, ERK, IKK α / β , caspase-3, and Ki-67 in colonic tissue. After a 7-day BCP treatment, colon samples were processed for immunohistochemical analysis. Preventive treatment with BCP (50 mg/kg, p.o.) significantly reduced phospho-p65, a nuclear factor B subunit (p65 NF κ B) (A), phospho-cAMP response element-binding (pCREB) (B), phospho-extracellular signal-regulated kinase (pERK) (C), phospho-I κ B-kinase α / β (pIKK α / β) (D), cleaved caspase-3 (E), and Ki-67 (F) immunostaining. The mean intensity of p65 NF κ B, pCREB, pERK, pIKK α / β , cleaved caspase-3, and Ki-67 staining were determined from image analysis and are represented as arbitrary units. Data are reported as means $\pm$ SEM ($n = 5$ to 7 mice per group). * $P < 0.05$ versus the DSS-treated group; ** $P < 0.05$ versus the control group.

A–D); however, AM630 notably abolished the protective effect of BCP against macroscopic colon damage, colon length reduction and MPO activity (Figure 6, A–D).

We investigated how the activation of the CB2 might be implicated in the beneficial effect of BCP in LPS-stimulated IEC-6 cells. The *in vitro* stimulation of IEC-6 cells with LPS for 4 hours increased mRNA for TNF- α and cytokine CINC-1 levels, which were reduced by preincubation with BCP (10 μ mol/L) (Figure 6, F and G). Treatment with AM630 alone did not alter the cytokine levels observed in the LPS-treated group; however, AM630 notably abolished the anti-inflammatory effect of BCP, indicating a functional CB2-dependent mechanism in the production of inflammatory mediators.

Some recent studies have shown a close interaction between the CB2 and PPAR γ .^{13,44} Indeed, PPAR γ agonists ameliorate experimental colitis,^{45–47} and PPAR γ expression is reduced in patients with ulcerative colitis.²⁰ To verify whether PPAR γ was also involved in colitis improvement after BCP treatment, the PPAR γ antagonist GW9662 (1 mg/kg, i.p.) was administered once a day from day 0 to day 7, alone or in combination with BCP. As was observed for CB2 antagonism, the PPAR γ antagonist significantly blocked the beneficial effect of BCP in DSS-induced colitis by partially reversing the DAI score, mac-

roscopic colon damage, and colon length and by restoring the activity of MPO (Figure 7, A–D). Furthermore, DSS-treated mice showed a reduction in mRNA expression for PPAR γ , but treatment with BCP partially restored PPAR γ expression (Figure 7E).

Discussion

IBDs are chronic debilitating diseases of the gastrointestinal tract that affect millions of people worldwide. Current therapies for IBDs remain inadequate. In recent years, much effort has gone into identifying promising new targets for IBD treatment. In this context, the use of herbal therapy or active compounds from plants constitutes an attractive approach for IBD treatment.^{48,49} BCP is a secondary metabolite found in many plant species,^{50,51} and it exhibits long-lasting oral and topical anti-inflammatory properties in different inflammatory models.^{49,52}

Corroborating a previous report, the present study demonstrated that BCP, given orally, consistently ameliorated the inflammatory signs associated with DSS-induced colitis, such as decreased body weight and colon length and increased colonic tissue damage,⁵³ in both therapeutic and preventive treatment. In all parameters analyzed, preventive treatment with BCP was more effective than therapeutic treatment at the highest dose (50 mg/kg, p.o.), but the significant difference was small, and both treatments are likely to be of use for clinical applications of BCP. In addition, in the present study the improvement in DSS-induced colitis by BCP oral treatment was associated with a direct reduction in colonic cytokines, chemokines, neutrophil migration, and macrophage infiltration and activation. There was also an increase in IL-4 levels and Foxp3 expression. Of great relevance, the present study clearly showed that BCP treatment markedly reduced activation of ERK1/2, p65 NF κ B, IKK α / β , CREB, caspase-3, and Ki-67. Notably, the beneficial effect of BCP was dependent on CB2 and PPAR γ activation. We also revealed a novel finding, that BCP pretreatment ameliorated oxazolone-induced colitis by decreasing weight loss and enhancing survival, suggesting that the anti-inflammatory effect of BCP is not dependent on the animal model used.

Both macrophages and neutrophils have pathogenic roles in animal models of IBD.^{9,54} Mononuclear cell infiltration was analyzed indirectly by measuring NAG activity. Colon tissues had a high number of resident mononuclear cells, which was significantly enhanced 7 days after DSS administration. Treatment with BCP prevented the increase in NAG activity, suggesting an impairment of mononuclear cell infiltration. BCP also decreases MPO activity in inflamed colons⁵³ and in rat paws treated with carrageenan⁴⁹ or LPS.⁵² The reduction in cellular infiltration observed in the present study was associated with a decrease in colon damage. It has been suggested that the inhibition of migratory mononuclear cells,⁵⁵ but not resident cells,⁵⁶ ameliorates colitis. Similarly, the infiltration of neutrophils into the colonic mucosa is believed to play a key role in mediating tissue damage and the clinical symptoms in experimental colitis.⁹

Figure 6. Pharmacological blockade of CB₂ receptor reverses BCP anti-inflammatory effects. Mice were given DSS (3%) for 5 days and were treated from day 0 to day 7 with BCP (50 mg/kg, p.o.) alone, with the CB₂ selective antagonist AM630 (AM; 10 mg/kg, i.p.) alone, or with AM630 (10 mg/kg, i.p., 30 minutes before) plus BCP (50 mg/kg, p.o.). Preventive oral treatment with BCP ameliorated all of the parameters analyzed, but treatment with AM630 plus BCP impaired the BCP improvement in DAI score (A), BCP macroscopic colon damage amelioration (B), and the BCP protective effect on colon length (C) and on MPO activity (D). At the end of 7 days, DSS enhanced mRNA expression for CB₂ on mouse colonic tissue and BCP reduced the mRNA expression of CB₂ (E). IEC-6 cells from rats were stimulated with LPS (1 μg/well) in the presence or absence of BCP (10 μmol/L per well) for 24 hours; the cell lysate was analyzed for TNF-α mRNA expression and secreted cytokines in the supernatant were analyzed for CINC-1 levels by enzyme-linked immunosorbent assay. BCP incubation reduced levels of TNF-α mRNA (F) and CINC-1 (G). The real-time PCR assay was performed in duplicate and *GAPDH* mRNA was used to normalize the relative amount of mRNA. Data are reported as means ± SEM (*n* = 5 to 7 mice per group). **P* < 0.05 versus the DSS-treated group; ***P* < 0.05 versus the control group; ****P* < 0.05 versus the BCP-treated group.

The pathway underlying the BCP protection of colon damage observed with both preventive and therapeutic treatment could be associated with diminished cleaved caspase-3 and Ki-67, which are fundamental proteins involved in apoptosis and cell proliferation, respectively.^{39,40} These events are associated with chronic inflammation and colon cancer and are increased under these conditions.⁵⁷ Furthermore, hyperproliferation and the abnormal distribution of replicating cells along the length of the colonic crypts are a marker of increased susceptibility to colorectal cancer and chronic inflammation.⁵⁷ In the present study, BCP oral treatment significantly reduced caspase-3 and Ki-67 protein activation in colonic tissue, suggesting that BCP ameliorates acute colitis by reducing cell proliferation and apoptosis. However, these events might be an indirect consequence of a general improvement of inflammation by BCP. In addition, we observed that DSS-induced colitis was associated with significant decreases in the colonic expression of the tight junction protein claudin-4, as described previously by Oshima et al.⁵⁸ BCP treatment did not, however, restore the levels of this protein. Our data suggest that the anti-inflammatory effects of BCP do not rely on its ability to modify tight-junction protein expression, such as claudin-4.

The prevention of cell infiltration and colon damage observed in mice orally treated with BCP was also associated with reduced protein levels and mRNA expression

of TNF-α, IL-1β, INF-γ, and CXCL1/KC, which are soluble mediators involved in cellular migration and adhesion molecule up-regulation.³⁷ The decreased levels of colonic cytokines might therefore explain the reduction of cell infiltration; however, because inflammatory cells produce cytokines and chemokines, the reduction of these mediators in the colon could be simply a consequence of a decreased cell migration. To clarify this hypothesis, we performed a lymph node and bone marrow-derived macrophage culture. We observed that BCP reduced CXCL1/KC, TNF-α, and MIP-2 levels by LPS-induced inflammation and the mRNA expression of CXCL1/KC and TNFα. This indicates that BCP not only decreases cell migration but also modulates cell activation. This finding confirms and extends previous work showing that BCP inhibits IL-1β and TNF-α protein expression in human peripheral whole blood stimulated by LPS.²³

The transcription of several cytokines and chemokines are controlled by ERK-NFκB,⁵⁹ and this transcription factor is activated during experimental colitis.³⁵ We therefore investigated whether the effect of BCP in decreasing the inflammatory mediators was associated with the ERK-NFκB signaling pathway. Our data demonstrated a marked inhibition of the p65 NFκB subunit and ERK-1/2 in the colons of mice treated with BCP. These results are in agreement with previous studies, in which BCP inhibited NFκB activation in the footpad of rats injected with LPS⁵² and diminished LPS-induced ERK1/2 phos-

Figure 7. Pharmacological blockade of PPAR γ receptor partially reverses BCP anti-inflammatory effects on DSS-induced colitis. Mice were given DSS (3%) for 5 days and were treated from day 0 to day 7 with BCP (50 mg/kg, p.o.) alone, the PPAR γ selective antagonist GW9662 (1 mg/kg, i.p.) alone, or with GW9662 (1 mg/kg, i.p., 30 minutes before) plus BCP (50 mg/kg, p.o.). Preventive oral treatment with BCP ameliorated all clinical parameters analyzed, but treatment with GW9662 plus BCP partially reduced BCP improvement in DAI score (A), BCP macroscopic colon damage amelioration (B), and BCP protective effect on colon length (C) and on MPO activity (D). At the end of 7 days, DSS diminished mRNA expression for PPAR γ in the mouse colon and BCP partially recovered the mRNA expression of PPAR γ (E). Real-time PCR assay was performed in duplicate; GAPDH mRNA was used to normalize the relative amount of mRNA. Data are reported as means $\pm$ SEM ($n = 5$ to 7 mice per group). * $P < 0.05$ versus the DSS-treated group; ** $P < 0.05$ versus the control group; *** $P < 0.05$ versus the BCP-treated group.

phorylation in human monocytes.²³ Moreover, a strong activation of IKK α/β was observed in DSS-treated mice, and this effect was suppressed by BCP treatment. Eckmann et al⁶⁰ showed that IKK α/β is closely involved in the proinflammatory response in acute colitis induced by DSS, suggesting that blockade of this kinase may be an important target for decreasing the inflammatory response.

Another transcription factor known to be involved in experimental colitis is CREB.³⁵ Genes whose transcription is regulated by CREB include those encoding the T helper type 1 (T_H1) cytokines.^{61,62} The T_H1 cytokines encompass the cytokine profile of acute murine DSS-induced colitis (INF- γ , TNF- α , and IL-1 β).⁶³ In T_H1 cells, the recruitment of phospho-CREB is significantly higher than in T_H2 cells.⁶⁴ In the present study, we demonstrated that BCP treatment decreased DSS-induced CREB activation, which may be linked to decreased T_H1 cytokines, such as INF- γ , and the enhancement of T_H2 cytokines, namely IL-4. In addition, IL-4 is associated with alternative macrophage activation, polarizing macrophages to the anti-inflammatory phenotype (M2).⁶⁵ Of note, *in vitro* macrophage treatment with BCP decreased proinflammatory cytokine production.

T_H2 regulatory T cells can also produce IL-4.⁶⁶ The present study demonstrated that BCP treatment enhanced the expression of FoxP3, the most reliable regulatory T-cell marker. Therefore, those cells could also be the source of IL-4. Nonetheless, regulatory T cells are known to produce transforming growth factor- β and IL-10, anti-inflammatory cytokines that were not altered by BCP treatment. The role of these cells in amelioration of BCP-induced colitis should be further examined.

Polarization toward T_H2 immunity can also be achieved by cannabinoids in a variety of models. Recently, Gertsch et al²³ reported that BCP is a CB2 agonist. CB2 is expressed mainly in immune cells, and its activation limits intestinal inflammation³⁰ and reduces proinflammatory mediator release.⁶⁷ The present study demonstrated that the anti-inflammatory effect of BCP was almost completely reversed by the selective CB2 antagonist AM630, strongly suggesting that a great part of the anti-inflammatory properties of BCP observed in DSS-induced colitis was mediated by CB2 activation. One of the possible mechanisms of action of BCP is the activation of CB2s and the consequent regulation of the activation and balance of T_H1/T_H2 cells. To confirm this hypothesis, we

performed another experiment in which we measured both the levels and expression of some cytokines *in vitro*. Our results showed that the incubation with AM630 reversed the inhibitory effect of BCP on levels of proinflammatory chemokine CINC-1/CXCL1 (cytokine chemoattractant for neutrophils) and on TNF- α mRNA expression in the IEC-6 cell line.

We performed another experiment to explore further whether BCP triggers CB2-dependent effects on cytokine levels/expression *in vitro*. Our results showed that the incubation with the CB2 antagonist AM630 reversed the inhibitory effect of BCP on proinflammatory chemokine CINC-1/CXCL1 levels and TNF- α mRNA expression in the IEC-6 cell line. These results are in agreement with Gertsch et al,²³ who found that AM630 reversed the inhibitory effect of BCP on IL-1 β and TNF- α protein expression in human peripheral whole blood. The present results provide convincing experimental evidence indicating that the effects of BCP depend on CB2 activation.

A shift toward a T_H2 cytokine dominance has been observed also in DSS-induced inflammation after PPAR γ ligand treatment. The PPAR γ ligands troglitazone, pioglitazone, and rosiglitazone decrease INF- γ and TNF- α levels and increase IL-4 levels and GATA-3 expression.⁶⁸ Other evidence supports a role of the PPAR γ in the regulation of intestinal inflammation: i) there is an impaired PPAR γ expression in colonic epithelial cells of patients with ulcerative colitis²⁰; ii) PPAR γ agonists dramatically reduce disease severity in mice²²; and iii) an increased susceptibility to colitis is observed in PPAR γ heterozygous mice (PPAR $\gamma^{+/-}$).²² A novel finding in the present study was that the anti-inflammatory effects of BCP were markedly reduced by the PPAR γ selective antagonist GW9662, demonstrating an important role of this nuclear transcription factor in the mechanism of the anti-inflammatory action of BCP.

Some evidence suggests cross-talk between PPAR γ and the cannabinoid pathways. Liu et al¹³ demonstrated the activation and binding of PPAR γ by synthetic cannabinoid. O'Sullivan et al¹⁴ demonstrated that Δ^9 -tetrahydrocannabinol causes vasorelaxation through activation of PPAR γ . Both CB2 and PPAR γ agonists reduce MPO activity^{29,69} and increase a T_H2 cytokine profile^{68,70} in experimental colitis in mice. In addition, the activation of CB2 and/or PPAR γ significantly reduces CXCL1/KC and MIP-2 tissue levels^{71,72} and inhibits NF κ B, IKK α/β , and CREB activation.^{45,73–75} As we have noted, all of these effects were observed in the colons from mice treated with BCP. Collectively, therefore, these data reinforce the important role of CB2 and PPAR γ activation in BCP anti-inflammatory activity.

Although BCP was suggested to act mainly on CB2, high doses of BCP induce anti-edematogenic effects in CB2 knockout mice,²³ indicating that BCP may not act exclusively on this receptor. In fact, some CB2 agonists can directly bind and activate PPAR γ .^{7,13,76} It is noteworthy that BCP is a lipophilic molecule that easily crosses the cell membrane²³ and that could, therefore, exert its pharmacological effects by acting directly on intracellular pathways. Indeed, the exact mechanisms by which cannabinoids interact with PPARs remain unclear.

Three potential pathways have been suggested to be involved in the mechanism of the activation of PPARs by cannabinoids: i) direct binding to PPARs; ii) the conversion of cannabinoids into metabolites that in turn activate PPARs; and iii) the interaction of cannabinoids with cell surface receptors, initiating intracellular signaling cascades that lead to the activation of PPARs. The present study demonstrated that the beneficial effects of BCP on DAI score were consistently reversed by CB2 and PPAR γ antagonists. Therefore, CB2 activation by BCP may trigger PPAR γ activation. We cannot, however, dismiss the possibility of direct action of BCP on the PPAR γ pathway. Furthermore, the fact that agonists of both CB2 and PPAR γ significantly improved experimental colitis could not be directly related with their expression in inflamed tissue, but it is not fully understood why mRNA levels of CB2 is overexpressed but mRNA levels of PPAR γ are underexpressed during DSS-induced colitis.

In conclusion, the present results demonstrate that the plant-derived sesquiterpene BCP, given orally, exhibits both preventive and therapeutic effects in DSS-induced colitis in mice. The mechanisms were associated with activation of CB2 and PPAR γ receptors, leading to the inhibition of proinflammatory cytokines and inflammatory cell influx. All of these effects were also associated with the inhibition of nuclear factors NF κ B and CREB and a shift toward T_H2 cytokines. Taken together, the present findings strongly suggest that BCP could constitute an attractive and apparently safe molecule for development of new anti-inflammatory drugs with therapeutic potential for use in treatment of human IBDs, such as ulcerative colitis and Crohn's disease.

Acknowledgment

We thank Prof. Dr. Patrícia Bozza for providing us with the IEC-6 cell line.

References

1. Cuzzocrea S: Emerging biotherapies for inflammatory bowel disease. *Expert Opin Emerg Drugs* 2003, 8:339–347
2. Podolsky DK: The current future understanding of inflammatory bowel disease. *Best Pract Res Clin Gastroenterol* 2002, 16:933–943
3. McGuckin MA, Eri R, Simms LA, Florin TH, Radford-Smith G: Intestinal barrier dysfunction in inflammatory bowel diseases. *Inflamm Bowel Dis* 2009, 15:100–113
4. Roberds SL, Anderson J, Basi G, Bienkowski MJ, Branstetter DG, Chen KS, Freedman SB, Frigon NL, Games D, Hu K, Johnson-Wood K, Kappenman KE, Kawabe TT, Kola I, Kuehn R, Lee M, Liu W, Motter R, Nichols NF, Power M, Robertson DW, Schenk D, Schoor M, Shopp GM, Shuck ME, Sinha S, Svensson KA, Tatsuno G, Tintrup H, Wijsman J, Wright S, McConlogue L: BACE knockout mice are healthy despite lacking the primary beta-secretase activity in brain: implications for Alzheimer's disease therapeutics. *Hum Mol Genet* 2001, 10:1317–1324
5. D'Argenio G, Valenti M, Scaglione G, Cosenza V, Sorrentini I, Di Marzo V: Up-regulation of anandamide levels as an endogenous mechanism and a pharmacological strategy to limit colon inflammation. *FASEB J* 2006, 20:568–570
6. Engel MA, Kellermann CA, Burnat G, Hahn EG, Rau T, Konturek PC: Mice lacking cannabinoid CB1-, CB2-receptors or both receptors show increased susceptibility to trinitrobenzene sulfonic acid (TNBS)-induced colitis. *J Physiol Pharmacol* 61:89–97

7. Galiègue S, Mary S, Marchand J, Dussosoy D, Carrière D, Carayon P, Bouaboula M, Shire D, Le Fur G, Casellas P: Expression of central and peripheral cannabinoid receptors in human immune tissues and leukocyte subpopulations. *Eur J Biochem* 1995, 232:54–61
8. Maresz K, Pryce G, Ponomarev ED, Marsicano G, Croxford JL, Shriver LP, Ledent C, Cheng X, Carrier EJ, Mann MK, Giovannoni G, Pertwee RG, Yamamura T, Buckley NE, Hillard CJ, Lutz B, Baker D, Dittel BN: Direct suppression of CNS autoimmune inflammation via the cannabinoid receptor CB1 on neurons and CB2 on autoreactive T cells. *Nat Med* 2007, 13:492–497
9. Bento AF, Leite DF, Claudino RF, Hara DB, Leal PC, Calixto JB: The selective nonpeptide CXCR2 antagonist SB225002 ameliorates acute experimental colitis in mice. *J Leukoc Biol* 2008, 84:1213–1221
10. Kühl AA, Kakirman H, Janotta M, Dreher S, Cremer P, Pawlowski NN, Loddenkemper C, Heimesaat MM, Grollich K, Zeitz M, Farkas S, Hoffmann JC: Aggravation of different types of experimental colitis by depletion or adhesion blockade of neutrophils. *Gastroenterology* 2007, 133:1882–1892
11. Morgan NH, Stanford IM, Woodhall GL: Functional CB2 type cannabinoid receptors at CNS synapses. *Neuropharmacology* 2009, 57:356–368
12. Van Sickle MD, Duncan M, Kingsley PJ, Mouihate A, Urbani P, Mackie K, Stella N, Makriyannis A, Piomelli D, Davison JS, Marnett LJ, Di Marzo V, Pittman QJ, Patel KD, Sharkey KA: Identification and functional characterization of brainstem cannabinoid CB2 receptors. *Science* 2005, 310:329–332
13. Liu J, Li H, Burstein SH, Zurier RB, Chen JD: Activation and binding of peroxisome proliferator-activated receptor gamma by synthetic cannabinoid ajulemic acid. *Mol Pharmacol* 2003, 63:983–992
14. O'Sullivan SE, Tarling EJ, Bennett AJ, Kendall DA, Randall MD: Novel time-dependent vascular actions of Delta9-tetrahydrocannabinol mediated by peroxisome proliferator-activated receptor gamma. *Biochem Biophys Res Commun* 2005, 337:824–831
15. Rousseaux C, Lefebvre B, Dubuquoy L, Lefebvre P, Romano O, Auwerx J, Metzger D, Wahli W, Desvergne B, Naccari GC, Chavatte P, Farce A, Bulois P, Cortot A, Colombel JF, Desreumaux P: Intestinal antiinflammatory effect of 5-aminosalicylic acid is dependent on peroxisome proliferator-activated receptor-gamma. *J Exp Med* 2005, 201:1205–1215
16. Schaefer KL, Denevich S, Ma C, Cooley SR, Nakajima A, Wada K, Schleizinger J, Sherr D, Saubermann LJ: Intestinal antiinflammatory effects of thiazolidenedione peroxisome proliferator-activated receptor-gamma ligands on T helper type 1 chemokine regulation include nontranscriptional control mechanisms. *Inflamm Bowel Dis* 2005, 11:244–252
17. Jiang C, Ting AT, Seed B: PPAR-gamma agonists inhibit production of monocyte inflammatory cytokines. *Nature* 1998, 391:82–86
18. Ricote M, Li AC, Willson TM, Kelly CJ, Glass CK: The peroxisome proliferator-activated receptor-gamma is a negative regulator of macrophage activation. *Nature* 1998, 391:79–82
19. Wahli W: A gut feeling of the PXR, PPAR and NF-kappaB connection. *J Intern Med* 2008, 263:613–619
20. Dubuquoy L, Jansson EA, Deeb S, Rakotobe S, Karoui M, Colombel JF, Auwerx J, Pettersson S, Desreumaux P: Impaired expression of peroxisome proliferator-activated receptor gamma in ulcerative colitis. *Gastroenterology* 2003, 124:1265–1276
21. Lefebvre M, Paulweber B, Fajas L, Woods J, McCrory C, Colombel JF, Najib J, Fruchart JC, Datz C, Vidal H, Desreumaux P, Auwerx J: Peroxisome proliferator-activated receptor gamma is induced during differentiation of colon epithelium cells. *J Endocrinol* 1999, 162:331–340
22. Buisine MP, Desreumaux P, Leteurtre E, Copin MC, Colombel JF, Porchet N, Aubert JP: Mucin gene expression in intestinal epithelial cells in Crohn's disease. *Gut* 2001, 49:544–551
23. Gertsch J, Leonti M, Raduner S, Racz I, Chen JZ, Xie XQ, Altmann KH, Karsak M, Zimmer A: Beta-caryophyllene is a dietary cannabinoid. *Proc Natl Acad Sci USA* 2008, 105:9099–9104
24. Gertsch J: Anti-inflammatory cannabinoids in diet: towards a better understanding of CB(2) receptor action? *Commun Integr Biol* 2008, 1:26–28
25. Orav A, Stulova I, Kailas T, Müürisepp M: Effect of storage on the essential oil composition of *Piper nigrum* L. fruits of different ripening states. *J Agric Food Chem* 2004, 52:2582–2586
26. de Carvalho PM Jr, Rodrigues RF, Sawaya AC, Marques MO, Shimizu MT: Chemical composition and antimicrobial activity of the essential oil of *Cordia verbenacea* D.C. *J Ethnopharmacol* 2004, 95:297–301
27. Ghia JE, Blennerhassett P, Collins SM: Impaired parasympathetic function increases susceptibility to inflammatory bowel disease in a mouse model of depression. *J Clin Invest* 2008, 118:2209–2218
28. Vandembark AA, Chou YK, Whitham R, Mass M, Buenafe A, Liefeld D, Kavanagh D, Cooper S, Hashim GA, Offner H: Treatment of multiple sclerosis with T-cell receptor peptides: results of a double-blind pilot trial [Erratum appeared in *Nat Med* 1997;3:240]. *Nat Med* 1996, 2:1109–1115
29. Kimball ES, Wallace NH, Schneider CR, D'Andrea MR, Hornby PJ: Vanilloid receptor 1 antagonists attenuate disease severity in dextran sulphate sodium-induced colitis in mice. *Neurogastroenterol Motil* 2004, 16:811–818
30. Storr MA, Keenan CM, Zhang H, Patel KD, Makriyannis A, Sharkey KA: Activation of the cannabinoid 2 receptor (CB2) protects against experimental colitis. *Inflamm Bowel Dis* 2009, 15:1678–1685
31. Napimoga MH, Vieira SM, Dal-Secco D, Freitas A, Souto FO, Mestriner FL, Alves-Filho JC, Grespan R, Kawai T, Ferreira SH, Cunha FQ: Peroxisome proliferator-activated receptor-gamma ligand, 15-deoxy-Delta12,14-prostaglandin J2, reduces neutrophil migration via a nitric oxide pathway. *J Immunol* 2008, 180:609–617
32. Rogerio AP, Andrade EL, Leite DF, Figueiredo CP, Calixto JB: Preventive and therapeutic anti-inflammatory properties of the sesquiterpene alpha-humulene in experimental airways allergic inflammation. *Br J Pharmacol* 2009, 158:1074–1087
33. Bradford MM: A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* 1976, 72:248–254
34. Stanley ER: Murine bone marrow-derived macrophages. *Methods Mol Biol* 1997, 75:301–304
35. Vitor CE, Figueiredo CP, Hara DB, Bento AF, Mazzucco TL, Calixto JB: Therapeutic action and underlying mechanisms of a combination of two pentacyclic triterpenes, alpha- and beta-amyrin, in a mouse model of colitis. *Br J Pharmacol* 2009, 157:1034–1044
36. Baumgart DC, Carding SR: Inflammatory bowel disease: cause and immunobiology. *Lancet* 2007, 369:1627–1640
37. Barker JN, Jones ML, Swenson CL, Sarma V, Mitra RS, Ward PA, Johnson KJ, Fantone JC, Dixit VM, Nickoloff BJ: Monocyte chemotaxis and activating factor production by keratinocytes in response to IFN-gamma. *J Immunol* 1991, 146:1192–1197
38. Arulampalam V, Pettersson S: Uncoupling the p38 MAPK kinase in IBD: a double edged sword? *Gut* 2002, 50:446–447
39. Harrington HA, Ho KL, Ghosh S, Tung KC: Construction and analysis of a modular model of caspase activation in apoptosis. *Theor Biol Med Model* 2008, 5:26
40. Gerdes J, Lemke H, Baisch H, Wacker HH, Schwab U, Stein H: Cell cycle analysis of a cell proliferation-associated human nuclear antigen defined by the monoclonal antibody Ki-67. *J Immunol* 1984, 133:1710–1715
41. Tsukita S, Furuse M: The structure and function of claudins, cell adhesion molecules at tight junctions. *Ann N Y Acad Sci* 2000, 915:129–135
42. Klein TW: Cannabinoid-based drugs as anti-inflammatory therapeutics. *Nat Rev Immunol* 2005, 5:400–411
43. Massa F, Marsicano G, Hermann H, Cannich A, Monory K, Cravatt BF, Ferri GL, Sibaev A, Storr M, Lutz B: The endogenous cannabinoid system protects against colonic inflammation. *J Clin Invest* 2004, 113:1202–1209
44. Burstein S: PPAR-gamma: a nuclear receptor with affinity for cannabinoids. *Life Sci* 2005, 77:1674–1684
45. Su CG, Wen X, Bailey ST, Jiang W, Rangwala SM, Keilbaugh SA, Flanagan A, Murthy S, Lazar MA, Wu GD: A novel therapy for colitis utilizing PPAR-gamma ligands to inhibit the epithelial inflammatory response. *J Clin Invest* 1999, 104:383–389
46. Takagi T, Naito Y, Tomatsuri N, Hando O, Ichikawa H, Yoshida N, Yoshikawa T: Pioglitazone, a PPAR-gamma ligand, provides protection from dextran sulfate sodium-induced colitis in mice in association with inhibition of the NF-kappaB-cytokine cascade. *Redox Rep* 2002, 7:283–289
47. Sánchez-Hidalgo M, Martín AR, Villegas I, de la Lastra CA: Rosiglitazone, a PPARgamma ligand, modulates signal transduction path-

- ways during the development of acute TNBS-induced colitis in rats. *Eur J Pharmacol* 2007, 562:247–258
48. Fernandes ES, Passos GF, Campos MM, Araújo JG, Pesquero JL, Avellar MC, Teixeira MM, Calixto JB: Mechanisms underlying the modulatory action of platelet activating factor (PAF) on the upregulation of kinin B1 receptors in the rat paw. *Br J Pharmacol* 2003, 139:973–981
49. Passos GF, Fernandes ES, Campos MM, Araújo JG, Pesquero JL, Souza GE, Avellar MC, Teixeira MM, Calixto JB: Kinin B1 receptor up-regulation after lipopolysaccharide administration: role of proinflammatory cytokines and neutrophil influx. *J Immunol* 2004, 172:1839–1847
50. Kanokmedhakul S, Kanokmedhakul K, Lekphrom R: Bioactive constituents of the roots of *Polyalthia cerasoides*. *J Nat Prod* 2007, 70:1536–1538
51. Ogura Y, Bonen DK, Inohara N, Nicolae DL, Chen FF, Ramos R, Britton H, Moran T, Karaliuskas R, Duerr RH, Achkar JP, Brant SR, Bayless TM, Kirschner BS, Hanauer SB, Núñez G, Cho JH: A frame-shift mutation in NOD2 associated with susceptibility to Crohn's disease. *Nature* 2001, 411:603–606
52. Medeiros GF, Passos GF, Vitor CE, Koepp J, Mazzuco TL, Pianowski LF, Campos MM, Calixto JB: Effect of two active compounds obtained from the essential oil of *Cordia verbenacea* on the acute inflammatory responses elicited by LPS in the rat paw. *Br J Pharmacol* 2007, 151:618–627
53. Cho JY, Chang HJ, Lee SK, Kim HJ, Hwang JK, Chun HS: Amelioration of dextran sulfate sodium-induced colitis in mice by oral administration of beta-caryophyllene, a sesquiterpene. *Life Sci* 2007, 80:932–939
54. Ohkawara T, Nishihira J, Takeda H, Hige S, Kato M, Sugiyama T, Iwanaga T, Nakamura H, Mizue Y, Asaka M: Amelioration of dextran sulfate sodium-induced colitis by anti-macrophage migration inhibitory factor antibody in mice. *Gastroenterology* 2002, 123:256–270
55. Kanai T, Watanabe M, Okazawa A, Sato T, Yamazaki M, Okamoto S, Ishii H, Totsuka T, Iiyama R, Okamoto R, Ikeda M, Kurimoto M, Takeda K, Akira S, Hibi T: Macrophage-derived IL-18-mediated intestinal inflammation in the murine model of Crohn's disease. *Gastroenterology* 2001, 121:875–888
56. Qualls JE, Kaplan AM, van Rooijen N, Cohen DA: Suppression of experimental colitis by intestinal mononuclear phagocytes. *J Leukoc Biol* 2006, 80:802–815
57. Vetusch A, Latella G, Sfera R, Caprilli R, Gaudio E: Increased proliferation and apoptosis of colonic epithelial cells in dextran sulfate sodium-induced colitis in rats. *Dig Dis Sci* 2002, 47:1447–1457
58. Oshima T, Miwa H, Joh T: Changes in the expression of claudins in active ulcerative colitis. *J Gastroenterol Hepatol* 2008, 23 Suppl 2:S146–S150
59. Granata F, Frattini A, Loffredo S, Del Prete A, Sozzani S, Marone G, Triggiani M: Signaling events involved in cytokine and chemokine production induced by secretory phospholipase A2 in human lung macrophages. *Eur J Immunol* 2006, 36:1938–1950
60. Eckmann L, Nebelsiek T, Fingerle AA, Dann SM, Mages J, Lang R, Robine S, Kagnoff MF, Schmid RM, Karin M, Arkan MC, Greten FR: Opposing functions of IKKbeta during acute and chronic intestinal inflammation. *Proc Natl Acad Sci USA* 2008, 105:15058–15063
61. Cippitelli M, Sica A, Viggiano V, Ye J, Ghosh P, Birrer MJ, Young HA: Negative transcriptional regulation of the interferon-gamma promoter by glucocorticoids and dominant negative mutants of c-Jun. *J Biol Chem* 1995, 270:12548–12556
62. Samten B, Ghosh P, Yi AK, Weis SE, Lakey DL, Gonsky R, Pendurthi U, Wize B, Zhang Y, Zhang M, Gong J, Fernandez M, Safi H, Vankayalapati R, Young HA, Barnes PF: Reduced expression of nuclear cyclic adenosine 5'-monophosphate response element-binding proteins and IFN-gamma promoter function in disease due to an intracellular pathogen. *J Immunol* 2002, 168:3520–3526
63. Egger B, Bajaj-Elliott M, MacDonald TT, Inglin R, Eysselein VE, Buchler MW: Characterisation of acute murine dextran sodium sulphate colitis: cytokine profile and dose dependency. *Digestion* 2000, 62:240–248
64. Yano S, Ghosh P, Kusaba H, Buchholz M, Longo DL: Effect of promoter methylation on the regulation of IFN-gamma gene during in vitro differentiation of human peripheral blood T cells into a Th2 population. *J Immunol* 2003, 171:2510–2516
65. Mantovani A, Sica A, Sozzani S, Allavena P, Vecchi A, Locati M: The chemokine system in diverse forms of macrophage activation and polarization. *Trends Immunol* 2004, 25:677–686
66. Duchmann R, Zeitz M: T regulatory cell suppression of colitis: the role of TGF-beta. *Gut* 2006, 55:604–606
67. Correa F, Mestre L, Docagne F, Guaza C: Activation of cannabinoid CB2 receptor negatively regulates IL-12p40 production in murine macrophages: role of IL-10 and ERK1/2 kinase signaling. *Br J Pharmacol* 2005, 145:441–448
68. Saubermann LJ, Nakajima A, Wada K, Zhao S, Terauchi Y, Kadowaki T, Aburatani H, Matsushashi N, Nagai R, Blumberg RS: Peroxisome proliferator-activated receptor gamma agonist ligands stimulate a Th2 cytokine response and prevent acute colitis. *Inflamm Bowel Dis* 2002, 8:330–339
69. Ramakers JD, Verstege MI, Thuijls G, Te Velde AA, Mensink RP, Plat J: The PPARgamma agonist rosiglitazone impairs colonic inflammation in mice with experimental colitis. *J Clin Immunol* 2007, 27:275–283
70. Lu T, Newton C, Perkins I, Friedman H, Klein TW: Cannabinoid treatment suppresses the T-helper cell-polarizing function of mouse dendritic cells stimulated with *Legionella pneumophila* infection. *J Pharmacol Exp Ther* 2006, 319:269–276
71. Tschöp J, Kasten KR, Nogueiras R, Goetzman HS, Cave CM, England LG, Dattilo J, Lentsch AB, Tschöp MH, Caldwell CC: The cannabinoid receptor 2 is critical for the host response to sepsis. *J Immunol* 2009, 183:499–505
72. Tambe Y, Tsujiuchi H, Honda G, Ikeshiro Y, Tanaka S: Gastric cytoprotection of the non-steroidal anti-inflammatory sesquiterpene, beta-caryophyllene. *Planta Med* 1996, 62:469–470
73. Konstantinopoulos PA, Vondoros GP, Sotiropoulou-Bonikou G, Kominea A, Papavassiliou AG: NF-kappaB/PPAR gamma and/or AP-1/PPAR gamma 'on/off' switches and induction of CBP in colon adenocarcinomas: correlation with COX-2 expression. *Int J Colorectal Dis* 2007, 22:57–68
74. Herring AC, Kaminski NE: Cannabinol-mediated inhibition of nuclear factor-kappaB, cAMP response element-binding protein, and interleukin-2 secretion by activated thymocytes. *J Pharmacol Exp Ther* 1999, 291:1156–1163
75. Zhao CY, Wang YD, Zhou JY, Jia B, Cui JF: [An experimental study on the reverse mechanism of PPAR-gamma agonist rosiglitazone in rats with non-alcoholic steatohepatitis]. *Chinese. Zhonghua Gan Zang Bing Za Zhi* 2007, 15:450–455
76. O'Sullivan SE: Cannabinoids go nuclear: evidence for activation of peroxisome proliferator-activated receptors. *Br J Pharmacol* 2007, 152:576–582