c-mip Down-Regulates NF- κ B Activity and Promotes Apoptosis in Podocytes

Virginie Ory,^{*,†} Qingfeng Fan,^{*,†}
Nabila Hamdaoui,^{*,†} Shao-yu Zhang,^{*,†}
Dominique Desvaux,^{*,†‡} Vincent Audard,^{*,†}
Marina Candelier,^{*,†} Laure-Helene Noel,[§]
Philippe Lang,^{*,†¶||} Georges Guellaën,^{*,†}
Andre Pawlak,^{*,†} and Djillali Sahali^{*,†¶||}

From INSERM,* U955; University of Paris Faculty of Medicine[†]
and French Institute for Research in Nephrology and
Transplantation,^{||} UMRS 955, Equipe 21; the Service of Histology,
Department of Pathology,[¶] and the Service of Nephrology,[¶]
AP-HP, Group Hospital Henri Mondor Albert Chenevier, Créteil;
and the Service of Anatomicpathology,[§] Hospital Necker and
the University of Paris Descartes, UMRS1016, Paris, France

The mechanisms of podocyte disorders in cases of idiopathic nephrotic syndrome (INS) are complex and remain incompletely elucidated. The abnormal regulation of NF- κ B may play a key role in the pathophysiology of these podocyte diseases, but at present, NF- κ B has not been thoroughly investigated. In this study, we report that induction of c-mip in podocytes of patients with INS is associated with a down-regulation of RelA, a potent antiapoptotic factor that belongs to the NF- κ B family. Overexpression of c-mip in differentiated podocytes promotes apoptosis by inducing caspase-3 activity and up-regulating the proapoptotic protein Bax, whereas the overall levels of the antiapoptotic protein Bcl-2 was concomitantly decreased. The associated overexpression of RelA prevented the proapoptotic effects of c-mip. In addition, the targeted induction of c-mip in podocytes *in vivo* inhibited the expression of the RelA protein and increased the Bax/Bcl-2 ratio. The expression of both c-mip and active caspase-3 increased in focal and segmental glomerulosclerosis biopsies, and both proteins displayed a close spatial relationship. These results suggest that alterations in NF- κ B activity might result from the up-regulation of c-mip and are likely to contribute to podocyte disorders in cases of INS. (Am J Pathol 2012; 180:2284–2292; <http://dx.doi.org/10.1016/j.ajpath.2012.02.008>)

Idiopathic nephrotic syndrome (INS) defines several entities, including minimal change nephrotic syndrome (MCNS)

and focal and segmental glomerulosclerosis (FSGS), which are considered to be prototypic podocyte diseases.¹ Podocytes are terminally differentiated cells that line the outer aspect of the glomerular basement membrane and constitute the ultimate barrier to urinary protein loss by the formation and maintenance of the podocyte foot processes and the interposed slit diaphragm.² Podocyte diseases might result from genetic defects of proteins playing a key structural and/or regulatory role in the integrity of the glomerular filtration barrier.³ Regardless of the underlying cause, the early podocyte damages are characterized by ultrastructural alterations of the slit diaphragm. Cellular injuries may progress through a reversible stage to podocyte depletion and glomerulosclerosis, which commonly complicate the outcome of chronic glomerular diseases in INS. In acquired INS, the mechanisms preceding these changes remain to be elucidated.

The NF- κ B family of transcription factors plays a central role in many cellular processes through the regulation of genes involved in immunity, inflammation, cell proliferation, differentiation, and apoptosis. In mammalian cells, the NF- κ B family consists of five members, NF- κ B1 (p105/p50), NF- κ B2 (p100/p52), RelA (p65), RelB, and cRel.⁴ The hallmark of the NF- κ B family is the presence in all members of a Rel-homology domain at the NH₂-terminus, responsible for DNA binding, dimerization, and association with the I- κ B inhibitory proteins.⁵ Among the NF- κ B proteins, only RelA, RelB, and c-Rel have a COOH-terminus transactivation domain, which is lacking in p50 and p52, so that RelA, RelB, and c-Rel homo- or heterodimers function as transcriptional activators,

Supported by an Avenir Program from INSERM, a grant from the French Kidney Foundation, Association pour l'Utilisation du Rein Artificiel (AURA), Assistance Publique des Hôpitaux de Paris (Programme hospitalier de recherche clinique), and a grant from the Fondation pour la Recherche Médicale (Q.F.). V.O. was supported by grants from the Ministère de la recherche and from the Société de Néphrologie-Laboratoire Fresenius. N.H. was supported by a joint grant from the Association Française contre les Myopathies and the French Kidney Foundation.

Accepted for publication February 7, 2012.

V.O. and Q.F. contributed equally to this work.

Address reprint requests to Djillali Sahali, M.D., Ph.D., Unité de Recherches INSERM U 955, Hôpital Henri Mondor, F 94010 Créteil, France. E-mail: dil.sahali@inserm.fr.

whereas p50 and p52 homodimers function as repressors.⁴ Although in most cases, Rel proteins act as transcriptional activators, they may repress some target genes.^{6,7} Because of its implication in many biological processes and the adverse consequences of its dysregulation, the activity of NF- κ B is tightly regulated at multiple translational and posttranslational levels. The most common regulatory mechanism relies on the actions of the inhibitory I- κ B proteins.⁸ In resting cells, I- κ B protein binds to the nuclear localization site of the Rel-homology domain, preventing the nuclear translocation of NF- κ B protein and thereby its transcriptional activity. On activation, I- κ B protein is phosphorylated, which allows its ubiquitination and subsequent degradation by proteasome. We recently shown that c-mip interacts with RelA and prevents its dissociation from the NF- κ B/I- κ B α complexes, resulting in the inhibition of nuclear translocation of NF- κ B and stabilization of I- κ B α .⁹

In human and experimental glomerular diseases, several mediators regulated by NF- κ B such as cytokines and adhesion molecules have been reported to play pathogenic roles in inflammatory and proliferative glomerular diseases.¹⁰ In nonproliferative glomerular diseases characterized by heavy proteinuria, such as minimal change disease and membranous nephropathy, it has been suggested that the excess of proteins in the tubules and their subsequent degradation induce tubular expression of many proinflammatory genes through NF- κ B activation.¹¹

To our knowledge, the mechanisms of NF- κ B regulation in the podocytes are little known. We recently reported that c-mip is overproduced in the podocytes during glomerular damages and interferes with signaling pathways playing a key role in podocyte function. Whether c-mip influences NF- κ B activity has not been investigated. We report here that c-mip represses *in vitro* and *in vivo* NF- κ B activity by down-regulating the expression of RelA. This effect was only observed in podocytes, whereas tubular cells displayed a contrasting increase of NF- κ B activity. Moreover, c-mip increases the levels of Bax and enhances caspase-3 activity, but it reduces the Bcl-2 level, suggesting that c-mip exerts a proapoptotic function.

Materials and Methods

Patients

The diagnosis of kidney disease was performed by renal biopsy, which was performed before the start of treatment. All patients had proteinuria above 3 g in 24 hours and severe hypoalbuminemia at the time of blood sampling. MCNS and FSGS were clinically classified as idiopathic in all cases.

Control for c-mip screening includes adult patients with glomerular diseases (IgA, lupus nephritis) who exhibited a nephrotic syndrome. Normal renal samples were supplied by the hospital tissue bank (Platform of Biological Resources, Henri Mondor Hospital) from patients undergoing nephrectomy for polar kidney tumor.

Plasmid Constructs, Reverse Transcription, and Quantitative PCR

The expression plasmids for c-mip and RelA have been previously described.^{12,13} Mouse total RNA was prepared from glomerular fractions isolated by graded sieving, using RNeasy kit (Qiagen, Courtaboeuf, France). Quantitative real-time RT-PCR (RT-qPCR) for RelA was performed using the oligonucleotides sense (5'-TGTTAC-CATCAGGGCAGATC-3') and anti-sense (5'-CAGGG-TACTCCATCAGCATG-3'). The samples (2 μ L of the reverse transcription reaction mixture, corresponding to 20 ng of total RNA) were amplified in a 20- μ L reaction mixture containing 0.5 mmol/L of each primer and 1 $\times$ Light-Cycler DNA Master SYBR Green buffer (Roche Molecular Biochemical, Mannheim, Germany). RT-qPCR conditions include an initial denaturing step at 95°C for 15 minutes, followed by 40 cycles (denaturing: 95°C, 15 seconds; annealing: 59°C, 30 seconds; extension, 72°C, 35 seconds).

Generation of c-mip Stable Podocyte Cell Lines

Conditionally immortalized mouse podocytes have been described elsewhere.¹⁴ Before transfection, podocytes were maintained at 60% confluence, under permissive conditions (cells were cultured in RPMI 1640 medium containing 10% fetal calf serum, 100 U/mL penicillin, 100 μ g/mL streptomycin, 50 U/mL γ -INF, at 33°C). The full-length human c-mip was inserted into a pDEST40 plasmid (Invitrogen, Carlsbad, CA). Proliferating podocytes were transfected with c-mip expression plasmid (1 μ g/10⁶ cells) using the Amaxa system. Following transfection, cells were maintained in complete medium for 24 hours, then Geneticin (antibiotic G-418 sulfate; Invitrogen) was added at 400 μ g/mL (100% lethal to untransfected cells). Stably transfected cells were selected in serial passages using Geneticin, then stored in liquid nitrogen or expanded under permissive conditions in the presence of Geneticin until they were in sufficient number. Podocyte clones were isolated by limiting dilution and stored in liquid nitrogen. For transient transfection, podocytes were transfected with c-mip expression plasmid (pDEST40) alone or cotransfected with empty vector or RelA, using the Nanofectin method according to the instructions provided by the manufacturer (PAA Laboratories, Pasching, Austria). In some experiments, podocytes were incubated with 2 μ mol/L lactacystin (Sigma-Aldrich, St. Louis, MO) for 20 hours, and then protein lysates were prepared as previously described.¹³

Caspase-3 Activity Assays

Caspase-3 activity was monitored in a quantitative assay using a fluorogenic substrate. Podocytes stably transfected with c-mip or expressing empty vector (~200,000 cells) were grown on 60-mm dishes at 37°C for 96 hours. Floating cells were collected and centrifuged, and the pellet was lysed in 50 μ L of lysis buffer containing 50 mmol/L HEPES (pH 7.4), 100 mmol/L NaCl, 1% Nonidet P-40, 1 mmol/L EDTA (pH 8.0), 1 mmol/L dithiothreitol, 2

$\mu\text{g/mL}$ leupeptin, 2 $\mu\text{g/mL}$ aprotinin. Adherent cells were washed three times with cold PBS and lysed for 10 minutes on ice in 0.2 mL of lysis buffer. The lysates from adherent and floating cells were pooled and centrifuged, and the supernatants were collected. The activity of caspase recognizing the DEVD (Asp-Glu-Val-Asp) motif (DEVDase activity) was measured in 200 μL of assay buffer containing 100 mmol/L HEPES (pH 7.4), 10% sucrose, 10 mmol/L dithiothreitol, 500 $\mu\text{mol/L}$ EDTA, 50 μg of protein, and 20 $\mu\text{mol/L}$ N-acetyl-Asp-Glu-Val-Asp-7-amino-4-trifluoromethyl coumarin (AC-DEVD-AFC) as fluorogenic substrate (Biomol, Tebu, France). After 3 hours at 37°C, the fluorescence of the reaction mixture was monitored every 30 minutes for 7 hours, using a spectrofluorometer (FL600 Fluorescence Microplate Reader; Biotek Instruments, Winooski, VT), with excitation and emission wavelengths of 400 and 530 nm, respectively.

TUNEL Assays

Paraffin-embedded kidney tissue sections were dewaxed, rehydrated, and then treated with proteinase K solution [10 $\mu\text{g/mL}$ in 10 mmol/L Tris-HCl (pH 7.4)] for 30 minutes at room temperature. The sections were rinsed twice with PBS and the In Situ Cell Death Detection Kit, POD (Roche Diagnostics, Mannheim, Germany) was used to label apoptotic cells. The sections were incubated with 50 μL of TUNEL (terminal deoxynucleotidyl transferase biotin-dUTP nick end labeling) reaction mixture for 60 minutes at 37°C in a humidified atmosphere in the dark. Terminal transferase was omitted for the negative control. As a positive control, one section was incubated with DNase I [2000 U/mL in 50 mmol/L Tris-HCl (pH 7.5), 10 mmol/L MgCl_2 , and 1 mg/mL BSA] for 10 minutes at room temperature to induce DNA strand breaks, prior to the labeling procedures. The sections were rinsed three times with PBS, and then incubated with 50 μL of Converter-POD on the sample, for 30 minutes at 37°C. The sections were rinsed three times with PBS, and then 50 μL of diaminobenzidine substrate and incubated for several minutes at room temperature. Finally, the sections were rinsed with PBS, mounted under glass coverslips, and analyzed under a light microscope.

Light Microscopy and Immunohistochemistry Analysis

For light microscopy, the kidney sections from wild-type and c-mip transgenic (Tg) mice were incubated for 16 hours in Duboscq-Brasil solution, and subsequently dehydrated, paraffin embedded, and stained with periodic acid-Schiff reagent (PAS). For immunohistochemistry study, antigen retrieval was performed by immersing the slides in boiling 0.01 mol/L citrate buffer in a 500-W microwave oven for 15 minutes. The endogenous peroxidase activity was blocked with 0.3% H_2O_2 in methanol for 30 minutes. Slides were incubated with the blocking reagents consisting of the avidin-biotin solution for 30 minutes and the normal blocking serum for 20 minutes, and then incubated overnight with a specific polyclonal

antibody. After washing with PBS, they were incubated with biotinylated goat anti-rabbit antibody. An avidin-biotinylated horseradish peroxidase complex (Vectastain ABC Reagent; Vector Laboratories, Burlingame, CA) and 3,3'-diaminobenzidine (Sigma Biochemical, St. Louis, MO) as a chromogen were applied for visualization of the immunoreaction. Anti-RelA, anti-c-mip,¹³ and anti-cleaved caspase-3 (Asp175) were used at a final concentration of 1:50, 1:200, and 1:30 dilution, respectively. Omission of the primary antibody was considered a negative control.

Immunofluorescence

Immunofluorescence studies on kidney tissues were performed with 4- μm -thick cryostat sections fixed in acetone for 10 minutes, air-dried 30 minutes at room temperature, and then kept in PBS for 3 minutes and blocked in 1% BSA-PBS. The sections were incubated with the indicated antibodies for 1 hour at room temperature, washed with PBS, and incubated with fluorescein isothiocyanate-conjugated or red fluorescent dye-conjugated secondary antibodies. For double fluorochrome labeling, the slides were simultaneously incubated with rabbit anti-c-mip antibody and mouse anti-nephrin antibody. After washing with PBS, the slides were simultaneously incubated with fluorescein isothiocyanate-conjugated goat anti-rabbit IgG and cyanine 3-conjugated sheep anti-mouse IgG. Sections were examined by fluorescence microscopy (Carl Zeiss, Oberkochen, Germany) using red and green filters.

Antibodies and Western Blot Analysis

The primary antibodies used in this study included anti-RelA, anti-Tie2, anti-lamin B (Santa Cruz Biotechnology, Santa Cruz, CA), anti-cleaved caspase-3 (Asp 75), anti-Bax, (Cell Signaling Technology, Danvers, MA), anti-Bcl-2 (BD Transduction Laboratories; BD Biosciences, San Jose, CA), monoclonal anti-vimentin (Dako, Glostrup, Denmark), monoclonal anti-GAPDH (Abcam, Cambridge, UK), anti-actin (Sigma-Aldrich), and anti-nephrin (Progen, Heidelberg, Germany). The anti-c-mip polyclonal antibody has been previously described.¹⁵ Western blot analysis was performed, using classical methods.

Generation of c-mip Transgenic Mice

The generation of c-mip transgenic mice has been previously described.¹³ All experiments involving animals were conducted in accordance with French laws. All of the mice analyzed in this study were hemizygous males [Tg(+)] from the F_8 to F_{12} generations.

Histological Analysis

Morphological lesions were studied on 4-mm PAS-stained sections. Five kidney sections by mouse ($n = 5$ mice for each group) were analyzed. Between 20 and 30 glomeruli per kidney section were randomly selected.

Statistical Analysis

Statistical analyses of the data were performed using PRIZM 4 for Macintosh (GraphPad Software, La Jolla, CA). Unpaired or paired Student's *t*-tests were used. *P* values of less than 0.05 were considered significant.

Results

Expression of c-mip Is Associated With a Down-Regulation of NF- κ B in Podocytes of Patients with INS

Although the role of NF- κ B in podocytes is poorly understood, NF- κ B likely mediates important functions including regulation of signaling pathways and podocyte survival.¹⁶ To investigate the possible influence of c-mip on NF- κ B activity in podocytes of glomerular diseases, we studied the kidney biopsies of patients with INS. As previously reported, c-mip was clearly induced in kidney biopsies of patients with MCNS, but it was scarcely or not detected in normal human kidney.¹³ The distribution pattern shows that c-mip was not confined to a cytoplasmic compartment, it was also visualized in nuclei (Figure 1A). By contrast, we did not detect c-mip in glomeruli of IgA

Figure 1. Induction of c-mip in MCNS relapse is associated with a down-regulation of RelA. Representative immunohistochemical analysis of serial sections from normal human kidney (NHK) and kidney biopsy specimens. **A:** Expression of c-mip. Note that c-mip was below detection limits in normal glomeruli, whereas it was clearly visualized along the external side of the capillary loops of MCNS relapse. The expression of c-mip was not restricted to cytoplasm compartment but it is also distributed in the nuclei. Note that c-mip was not detected in IgA nephropathy or in proliferative lupus nephritis (LN-IV). Scale bars: 20 μ m. **B:** Expression of RelA in NHK and MCNS. Note that RelA is highly abundant in normal podocytes, whereas it was reduced in MCNS relapse. Scale bars: 20 μ m.

Figure 2. Confocal microscopy analysis. Detection of RelA (green); nephrin, tie-2, vimentin, and lamin-B (red) in normal human kidney (NHK, **left**) and in kidney biopsy specimens from MCNS relapse (**right**). Note that RelA partially colocalized with nephrin and lamin, but not with vimentin and tie-2 in normal glomeruli. The expression of RelA was hardly detected in MCNS relapse. Scale bars: 20 μ m.

nephropathy or proliferative lupus nephritis. Immunohistochemistry analysis on the same kidney specimens (5 normal specimens and 12 MCNS) using an anti-RelA antibody showed that RelA was clearly detected in normal podocytes, but it was dramatically altered in the podocytes of MCNS (Figure 1, A and B). Confocal microscopy analysis of normal glomeruli (Figure 2) shows that RelA colocalized partially with nephrin, but not with Tie2, suggesting that its expression is restricted to podocytes and did not overlap with vimentin but mostly colocalized with lamin-B, a marker of nuclear membrane. In some podocytes, RelA was also localized within the nucleus. By contrast, in glomeruli of patients with MCNS, the

Figure 3. The RelA protein is destabilized in the presence of c-mip, *in vivo* and *in vitro*. Glomeruli from wild-type (WT, $n = 5$) and 6-month-old c-mip transgenic mice [Tg(+), $n = 5$] were isolated, and the abundance of RelA was analyzed. **A:** Quantification of RelA transcripts by RT-qPCR. **B:** Quantitative Western blot analysis of RelA protein from WT and Tg(+) mice. Data are representative of three independent experiments. **C:** Podocytes were transiently transfected with c-mip or empty vector (Ev), cultured 20 hours with or without lactacystin (2 mmol/L), then analyzed by Western blotting. Note that lactacystin prevents the destabilization of endogenous RelA. Data are representative of two independent experiments.

abundance of RelA was lower (Figure 2). These results suggest that induction of c-mip in podocytes is associated with a down-regulation of RelA.

Expression of c-mip in Vivo Induces a Down-Regulation of RelA at the Posttranscriptional Level

To understand the functional consequences of c-mip induction in podocytes, we generated transgenic mice expressing c-mip under nephrin promoter to restrict transgene expression to podocytes. c-mip transgenic mice develop heavy proteinuria with foot process effacement without inflammatory lesions or immune complex deposits.¹³ The abundance of the RelA transcripts measured by quantitative RT-PCR did not significantly differ between wild-type and transgenic mice (Figure 3A). By contrast, Western blot analysis of glomerular protein lysates showed that RelA abundance was significantly decreased in c-mip transgenic mice, as compared with wild-type mice (Figure 3B), which suggests that c-mip affects only the stability of the RelA protein. Indeed, co-culture of c-mip-transfected podocytes in the presence of lactacystin, a potent proteasome inhibitor, prevented the decrease of endogenous RelA (Figure 3C), suggesting that c-mip targets RelA through proteasome-mediated degradation.

c-mip Displays Proapoptotic Properties

On the basis of the studies performed in NF- κ B-deficient mice, it is believed that NF- κ B is a potent antiapoptotic transcription factor.¹⁷ Given our findings showing a

down-regulation of RelA, which is essential for NF- κ B activity, we sought to determine whether c-mip influences the signaling pathways involved in apoptosis. Apoptosis is defined by morphological features including loss of adhesion, cell shrinkage, and biochemical alterations such as up-regulation of Bax and caspase-3 activity. Bax belongs to the Bcl-2 family of proteins, which include both antiapoptotic (eg, Bcl-2 and Bcl-xl) and proapoptotic (eg, Bax, Bad) members.¹⁸ Caspase-3 is one of the final downstream effectors of the apoptotic process, and its activation is closely associated with mitochondrial dysfunction.¹⁹ To assess whether c-mip is a proapoptotic molecule, we first transiently transfected c-mip alone in podocyte cell line and analyzed the expression level of Bax, BclII, and caspase-3. Overexpression of c-mip induced an up-regulation of Bax and caspase-3, whereas BclII abundance decreased when compared with empty vector-transfected cells (Figure 4A). On the other hand, cotransfection of RelA antagonized the proapoptotic effects of c-mip. We then sought to find whether these changes are reproducible in differentiated podocytes stably transfected with c-mip or empty vector. We studied five independent c-mip stably transfected podocyte cell lines and two control clones stably transfected with empty vector. We performed quantitative Western blot analysis to measure the abundance of the Bcl-2 and Bax proteins that were normalized to GAPDH loading. As shown in Figure 4B, Bax was overproduced in c-mip-overexpressing podocytes, so that the Bax/Bcl-2 ratio was significantly increased (Figure 4B). Morphological analyses showed that empty vector-transfected cells displayed small morphological changes, whereas c-mip-overexpressing podocytes exhibited severe cell shrinkage and most cells had detached. To assess whether

Figure 4. Transient and stable overexpression of c-mip in podocytes promotes apoptosis. **A:** Podocytes were singly transiently transfected with empty vector (Ev) or c-mip or cotransfected with Ev/c-mip or RelA/c-mip. The expression of Bax, BclII, and caspase-3 was analyzed by Western blotting on the protein lysates. Note that c-mip increases Bax and caspase-3 and down-regulates BclII, these alterations being prevented by the coexpression of RelA. **B:** Podocytes were stably transfected with c-mip expression plasmid (pDest 40) or empty vector (Ev), then cloned by limiting dilution. Morphological and/or quantitative Western blot data for six clones (five c-mip and one Ev clones) are shown. c-mip was revealed by anti-V5 tag monoclonal antibody. Quantification of c-mip was performed by subtracting the background related to the empty vector. Note that c-mip abundance varies depending on the clone; higher c-mip abundance is correlated with severe cellular injury and an increase of the Bax/Bcl-2 ratio. **C:** Differentiated podocytes stably transfected with c-mip or empty vector (Ev) were tested for caspase-3 activity assays. The caspase-3 activity was monitored every 30 minutes for 7 hours.

these morphological changes involve caspase-3 activation, we performed a caspase-3 activity assay on stable transfectant cells. Podocytes overexpressing c-mip exhibited an activation of caspase-3 in a time-dependent manner, whereas no significant change was detected in the empty vector-transfected podocytes (Figure 4C).

The relevance of these findings was analyzed in c-mip transgenic mice. In this model, mice develop after 3 months of age progressive FSGS.¹³ Histological analysis (Figure 5A) showed that the number of glomeruli displaying FSGS lesions was increased from 3 (4.0% $\pm$ 1.5%), 6 (7.5% $\pm$ 2.5%) to 12 months of age (23.0% $\pm$ 3.5%). Western blot analysis from glomerular extracts revealed increased Bax in c-mip Tg mice, whereas Bcl-2 was reduced, with a Bax/Bcl-2 ratio that was increased by 3.1 fold, as compared with wild-type mice (Figure 5B). We then used the TUNEL assay to detect DNA fragmentation in podocytes *in vivo*. In wild-type mice and in 3-month-old transgenic mice, TUNEL-positive nuclei were not detectable in glomeruli. However, in transgenic mice of 6 and 12 months of age, a number of podocytes became positive for TUNEL (Figure 6). Collectively, these results suggest that c-mip promotes apoptosis *in vitro* and *in vivo*.

FSGS lesions are characterized by dysregulation of podocyte function, leading to apoptosis. We have previously shown that c-mip abundance is increased in primary FSGS. Therefore, we investigated whether FSGS biopsy specimens exhibit active, cleaved caspase-3, a

marker of cell apoptosis. Immunocytochemistry analysis was performed in seven FSGS biopsy specimens. Active caspase-3 was detected in all examined cases, to a variable degree, along the external side of some capillary loops, consistent with podocyte localization (Figure 7). The expression was decreased or disappeared in podocyte-depleted areas corresponding to severe FSGS lesions or glomerulosclerosis (Figure 7). Interestingly, c-mip was increased in glomeruli in a pattern similar to that observed with active caspase-3.

Discussion

The role of NF- κ B in glomerular biology and in pathophysiology of podocyte disease remains little explored. In the present work, we provide evidence that i) increased expression of c-mip in podocytes of patients with INS is correlated with a down-regulation of RelA; ii) c-mip destabilizes endogenous RelA protein *in vitro*, which is prevented by lactacystin, suggesting that c-mip targets RelA through proteasome-mediated degradation; iii) stable transfection of c-mip in the podocytes is associated with cell shrinkage and increased abundance of Bax and active caspase-3; iv) c-mip induces *in vivo* an up-regulation of Bax and a decrease of Bcl-2, supporting the role of c-mip in apoptosis; v) c-mip transgenic mice develop progressive FSGS in the absence of overt tubular and

Figure 5. c-mip promotes progressive FSGS in transgenic mice. **A:** Representative Periodic acid-Schiff (PAS) staining of kidney sections from 3- to 12-month-old wild-type and proteinuric Tg mice. Scale bars: 50 μ m (top of each set); 20 μ m (bottom of each set). Glomerular alterations are apparent at 6 and 12 months, characterized by FSGS lesions and mesangial expansion. The tubules appear normal, and no inflammatory injuries can be seen. **Right,** percentage of glomeruli with FSGS. Five mice were analyzed by age range (≥ 300 glomeruli). Data represent the mean $\pm$ SEM. **B:** Glomerular protein extracts from 12-month-old wild-type and proteinuric Tg mice were analyzed by Western blotting for the relative expression of Bax and Bcl-2. Quantitative data for Bax/Bcl-2 ratio are shown to the right.

interstitial lesions; and vi) the abundance of c-mip and active caspase-3 was increased in human primary FSGS.

We have previously shown that c-mip interacts through its pleckstrin homology domain with Fyn and PI3 kinase regulatory subunit (p85), and exerts an inhibitory role in proximal signaling.^{13,20} c-mip also binds to and inhibits the activation of neuronal Wiskott-Aldrich syndrome protein (N-WASP), a major component of cytoskeleton that regulates the ARP2/3-mediated actin nucleation. In addition, we found that c-mip induces *in vitro* and *in vivo* cytoskeleton disorganization,¹³ and increases the abundance of death-associated protein (DAP) kinase.²⁰ DAP

kinase promotes apoptosis by disrupting integrin-mediated cell adhesion, alteration of cytoskeleton, and loss of extracellular matrix-dependent survival signals.²¹ Alteration of the actin network induces the production of reactive oxygen species from mitochondria, leading to apoptosis.²² We have previously found that c-mip binds RelA through its leucine-rich repeat-containing C-terminal domain.⁹ We show here that c-mip may target RelA to proteasome-dependent degradation. Altogether, our results suggest that c-mip is a multifunctional protein and may promote apoptosis through different mechanisms, including cytoskeleton alterations, up-regulation of DAP kinase, and down-regulation of RelA.

Several studies have shown that NF- κ B hyperactivation is mainly correlated with the intensity of proteinuria and is restricted in the tubules of patients with progressive kidney diseases, including glomerular diseases.^{23–25} Indeed, NF- κ B activity is associated with increased tubular expression of the chemokines monocyte chemoattractant protein-1 (MCP-1), regulated on activation normal T cell expressed and secreted (RANTES), osteopontin, and the profibrogenic cytokines platelet-derived growth factor-BB (PDGF-BB) and transforming growth factor- β (TGF β),²³ leading to tubular apoptosis and fibrosis. By contrast, these studies fail to detect significant hyperac-

Figure 6. c-mip promotes apoptosis *in vivo*. TUNEL assay on paraffin kidney sections of wild-type and c-mip transgenic mice at 3, 6, and 12 months of age. Note that TUNEL-positive cells were not detected in 3-month-old transgenic mice, whereas many podocytes became positive at 6 months of age. Scale bars: 20 μ m.

Figure 7. Expression of c-mip and caspase-3 in FSGS biopsies. Representative immunostaining of c-mip and cleaved caspase-3 from FSGS biopsy specimens.^{1–4} In most cases, the localization of c-mip and active caspase-3 was observed in close proximity. The staining strongly decreases or disappears in podocyte-depleted areas corresponding to FSGS lesions or severe glomerulosclerosis. Scale bars: 20 μ m.

tivation of NF- κ B in podocytes in certain forms of glomerular diseases, suggesting that this mechanism does not prevail in these cells. It has been reported that NF- κ B is expressed in a few mesangial cells and tubular epithelial cells in MCNS and membranous nephropathy, whereas it is detected in podocytes of IgA nephropathy and non-IgA mesangial proliferative glomerulonephritis.²⁴ These data are consistent with our results because the up-regulation of c-mip might account for the nondetection of NF- κ B in podocytes in INS. Conversely, in IgA nephropathy, where the expression of NF- κ B is found increased, c-mip was not detected in podocytes, in accord with previous data.¹³ The down-regulation of RelA in INS relapse may explain why these podocyte diseases are not associated with inflammatory processes in glomeruli. By contrast, activation of RelA in the podocytes of Par4-deficient mice is associated with severe glomerular and tubular inflammatory lesions.²⁶ Moreover, in pathological situations associated with sustained c-mip increase, the down-regulation of NF- κ B in glomeruli might represent a possible mechanism involved in podocyte apoptosis and may contribute to podocyte depletion in some podocyte diseases without early involvement of tubular and interstitial tissue injuries.

Two distinct signaling pathways are involved in the initiation of apoptosis: the extrinsic apoptotic pathway or receptor-linked pathway requires the binding of a ligand to a death receptor on the cell surface, such as TNF receptor; and the intrinsic apoptotic pathway that is mediated by mitochondrial damage and release of cytochrome *c*.²⁷ The development of the signaling cascade leading to apoptosis requires in most situations the up-regulation of p53. Despite several assays, we did not observe an up-regulation of p53 in stable transfected podocytes overexpressing c-mip or in c-mip Tg mice (data not shown), which suggests that the proapoptotic function of c-mip does not require p53. Some studies have shown that the ability to trigger apoptotic pathways in the absence of p53 may result from the activation of extracellular signal-regulated kinase (Erk) signaling.²⁸ However, the Erk pathway has multiple functions, including pro- and antiapoptotic effects, of which the mechanisms are still unclear.²⁹

Induction of apoptosis following exposition to puromycin aminonucleoside, in cultured podocytes as well as *in vivo*, in a model of FSGS in rats, is associated with an increase of Bax and a reduction of Bcl-2.^{19,30} The underlying mechanism involves an up-regulation of p53 since, in this model, apoptosis is prevented by pifithrin- α (PFT- α) and dexamethasone, which inhibit p53 activation. Indeed, two signaling pathways have been involved in podocyte apoptosis induced by puromycin aminonucleoside, depending on caspase-3 or apoptosis-inducing factor (AIF), respectively. Dexamethasone inactivates AIF by inhibiting its nuclear translocation.¹⁹

Insulin growth factor-1 (IGF-1) protects from apoptosis induced by etoposide, an inhibitor of topoisomerase II, through a mechanism involving activation of PI3 kinase and Bad.³¹ This conclusion is based on the fact that the inhibitors of PI3 kinase, Whormannin and Ly294002, abrogate the antiapoptotic effect of IGF1. Moreover, nephrin and vascular endothelial growth factor (VEGF) inhibit apoptosis by recruiting the PI3 kinase-Akt signaling pathway.³² Both the VEGF and nephrin signaling pathways appear interconnected because mutation of nephrin, which disrupts the nephrin signaling, abrogates the antiapoptotic effect of VEGF.³²

The antiapoptotic role of CD2-associated protein (CD2AP), an adapter protein anchoring the nephrin-mediated proximal signals to actin cytoskeleton, has been evidenced by the phenotype of CD2AP-deficient mice, which develop nephrotic syndrome and renal failure caused by glomerulosclerosis.³³ In this model, proteinuria occurs in the context of accelerated apoptosis resulting from hyperactivation of p38 mitogen-activated protein kinase (p38 MAPK) by TGF β 1.^{34,35} The mechanism by which CD2AP impedes the apoptotic process involves a down-regulation of p38 MAPK via activation of PI3 kinase and Akt signaling.³⁵

Direct evidence of the role of TGF β 1 in podocyte apoptosis comes from the study of transgenic mice overexpressing TGF β 1 under control of the albumin promoter.³⁶ In this model, Tg mice display glomerulosclerosis and interstitial fibrosis with renal failure, and death in 50% of offspring at 5 to 12 weeks of age.³⁴ Histological lesions

observed in TGF β 1 and c-mip Tg models are different. Glomerulosclerosis and extracellular matrix expansion are present in the early stages (~2 weeks) in TGF β 1 Tg mice, whereas glomeruli are morphologically normal despite the presence of a nephrotic syndrome and foot process fusions in c-mip Tg mice, until 4 to 5 months of age. Renal failure and death are rare during the first year of life in c-mip Tg mice even though these complications are common in TGF β 1 Tg mice. However, as with c-mip Tg mice, podocyte apoptosis precedes mesangial expansion.

In conclusion, we report here new functional aspects of c-mip in podocytes. We show that c-mip induces *in vitro* and *in vivo* the destabilization of the RelA protein in podocytes. The up-regulation of c-mip and active caspase-3 in human FSGS suggests that sustained up-regulation of c-mip might result in podocyte depletion and progressive FSGS.

Acknowledgments

We are grateful to Dr. Peter Mundel for providing us with the mouse podocyte cell line.

References

- Mathieson PW: Minimal change nephropathy and focal segmental glomerulosclerosis. *Semin Immunopathol* 2007, 29:415–426
- Tryggvason K, Patrakka J, Wartiovaara J: Hereditary proteinuria syndromes and mechanisms of proteinuria. *N Engl J Med* 2006, 354: 1387–1401
- Shankland SJ: The podocyte's response to injury: role in proteinuria and glomerulosclerosis. *Kidney Int* 2006, 69:2131–2147
- Shih VF, Tsui R, Caldwell A, Hoffmann A: A single NF-kappa B system for both canonical and non-canonical signaling. *Cell Res* 2011, 21: 86–102
- Zheng C, Yin Q, Wu H: Structural studies of NF-kappa B signaling. *Cell Res* 2011, 21:183–195
- Fu T, Li P, Wang H, He Y, Luo D, Zhang A, Tong W, Zhang L, Liu B, Hu C: c-Rel is a transcriptional repressor of EPHB2 in colorectal cancer. *J Pathol* 2009, 219:103–113
- Ashburner BP, Westerheide SD, Baldwin AS Jr.: The p65 (RelA) subunit of NF-kappa B interacts with the histone deacetylase (HDAC) corepressors HDAC1 and HDAC2 to negatively regulate gene expression. *Mol Cell Biol* 2001, 21:7065–7077
- Ferreiro DU, Komives EA: Molecular mechanisms of system control of NF-kappa B signaling by I-kappa B alpha. *Biochemistry* 2010, 49: 1560–1567
- Kamal M, Valanciute A, Dahan K, Ory V, Pawlak A, Lang P, Guellaen G, Sahali D: C-mip interacts physically with RelA and inhibits nuclear factor kappa B activity. *Mol Immunol* 2009, 46:991–998
- Mudge SJ, Paizis K, Auwardt RB, Thomas RJ, Power DA: Activation of nuclear factor-kappa B by podocytes in the autologous phase of passive Heymann nephritis. *Kidney Int* 2001, 59:923–931
- Remuzzi G, Ruggerenti P, Benigni A: Understanding the nature of renal disease progression. *Kidney Int* 1997, 51:2–15
- Valanciute A, le Gouvello S, Solhonne B, Pawlak A, Grimbort P, Lyonnet L, Hue S, Lang P, Remy P, Salomon R, Bensman A, Guellaen G, Sahali D: NF-kappa B p65 antagonizes IL-4 induction by c-maf in minimal change nephrotic syndrome. *J Immunol* 2004, 172:688–698
- Zhang SY, Kamal M, Dahan K, Pawlak A, Ory V, Desvaux D, Audard V, Candelier M, Mohamed FB, Matignon M, Christov C, Decroux X, Bernard V, Mangiapan G, Lang P, Guellaen G, Ronco P, Sahali D: c-mip impairs podocyte proximal signaling and induces heavy proteinuria. *Sci Signal* 2010, 3:ra39
- Mundel P, Reiser J, Zuniga Mejia Borja A, Pavenstadt H, Davidson GR, Kriz W, Zeller R: Rearrangements of the cytoskeleton and cell contacts induce process formation during differentiation of conditionally immortalized mouse podocyte cell lines. *Exp Cell Res* 1997, 236:248–258
- Audard V, Zhang SY, Copie-Bergman C, Rucker-Martin C, Ory V, Candelier M, Baia M, Lang P, Pawlak A, Sahali D: Occurrence of minimal change nephrotic syndrome in classical Hodgkin lymphoma is closely related to the induction of c-mip in Hodgkin-Reed Sternberg cells and podocytes. *Blood* 2010, 115:3756–3762
- Sanz AB, Sanchez-Nino MD, Ramos AM, Moreno JA, Santamaria B, Ruiz-Ortega M, Egido J, Ortiz A: NF-kappa B in renal inflammation. *J Am Soc Nephrol* 2010, 21:1254–1262
- Gerondakis S, Grumont R, Gugasyan R, Wong L, Isomura I, Ho W, Banerjee A: Unravelling the complexities of the NF-kappa B signalling pathway using mouse knockout and transgenic models. *Oncogene* 2006, 25:6781–6799
- Youle RJ, Strasser A: The BCL-2 protein family: opposing activities that mediate cell death. *Nat Rev Mol Cell Biol* 2008, 9:47–59
- Wada T, Pippin JW, Marshall CB, Griffin SV, Shankland SJ: Dexamethasone prevents podocyte apoptosis induced by puromycin aminonucleoside: role of p53 and Bcl-2-related family proteins. *J Am Soc Nephrol* 2005, 16:2615–2625
- Kamal M, Pawlak A, BenMohamed F, Valanciute A, Dahan K, Candelier M, Lang P, Guellaen G, Sahali D: C-mip interacts with the p85 subunit of PI3 kinase and exerts a dual effect on ERK signaling via the recruitment of Dp1 and DAP kinase. *FEBS Lett* 2010, 584:500–506
- Wang WJ, Kuo JC, Yao CC, Chen RH: DAP-kinase induces apoptosis by suppressing integrin activity and disrupting matrix survival signals. *J Cell Biol* 2002, 159:169–179
- Gourlay CW, Carpp LN, Timpson P, Winder SJ, Ayscough KR: A role for the actin cytoskeleton in cell death and aging in yeast. *J Cell Biol* 2004, 164:803–809
- Mezzano SA, Barria M, Droguett MA, Burgos ME, Ardiles LG, Flores C, Egido J: Tubular NF-kappa B and AP-1 activation in human proteinuric renal disease. *Kidney Int* 2001, 60:1366–1377
- Ashizawa M, Miyazaki M, Abe K, Furusu A, Isomoto H, Harada T, Ozono Y, Sakai H, Koji T, Kohno S: Detection of nuclear factor-kappa B in IgA nephropathy using Southwestern histochemistry. *Am J Kidney Dis* 2003, 42:76–86
- Zheng L, Sinniah R, Hsu SI: In situ glomerular expression of activated NF-kappa B in human lupus nephritis and other non-proliferative proteinuric glomerulopathy. *Virchows Arch* 2006, 448:172–183
- Hussain S, Romio L, Saleem M, Mathieson P, Serrano M, Moscat J, Diaz-Meco M, Scambler P, Kozliff A: Nephron deficiency activates NF-kappa B and promotes glomerular injury. *J Am Soc Nephrol* 2009, 20:1733–1743
- Reed JC: Mechanisms of apoptosis. *Am J Pathol* 2000, 157:1415–1430
- Tang D, Wu D, Hirao A, Lahti JM, Liu L, Mazza B, Kidd VJ, Mak TW, Ingram AJ: ERK activation mediates cell cycle arrest and apoptosis after DNA damage independently of p53. *J Biol Chem* 2002, 277: 12710–12717
- Wada T, Penninger JM: Mitogen-activated protein kinases in apoptosis regulation. *Oncogene* 2004, 23:2838–2849
- Wang W, Tzanidis A, Divjak M, Thomson NM, Stein-Oakley AN: Altered signaling and regulatory mechanisms of apoptosis in focal and segmental glomerulosclerosis. *J Am Soc Nephrol* 2001, 12:1422–1433
- Bridgewater DJ, Ho J, Sauro V, Matsell DG: Insulin-like growth factors inhibit podocyte apoptosis through the PI3 kinase pathway. *Kidney Int* 2005, 67:1308–1314
- Foster RR, Saleem MA, Mathieson PW, Bates DO, Harper SJ: Vascular endothelial growth factor and nephrin interact and reduce apoptosis in human podocytes. *Am J Physiol Renal Physiol* 2005, 288:F48–F57
- Shih NY, Li J, Karpitskii V, Nguyen A, Dustin ML, Kanagawa O, Miner JH, Shaw AS: Congenital nephrotic syndrome in mice lacking CD2-associated protein. *Science* 1999, 286:312–315
- Schiffer M, Bitzer M, Roberts IS, Kopp JB, ten Dijke P, Mundel P, Bottinger EP: Apoptosis in podocytes induced by TGF-beta and Smad7. *J Clin Invest* 2001, 108:807–816
- Schiffer M, Mundel P, Shaw AS, Bottinger EP: A novel role for the adaptor molecule CD2-associated protein in transforming growth factor-beta-induced apoptosis. *J Biol Chem* 2004, 279:37004–37012
- Kopp JB, Factor VM, Mozes M, Nagy P, Sanderson N, Bottinger EP, Klotman PE, Thorgeirsson SS: Transgenic mice with increased plasma levels of TGF-beta 1 develop progressive renal disease. *Lab Invest* 1996, 74:991–1003