GASTROINTESTINAL, HEPATOBILIARY, AND PANCREATIC PATHOLOGY

Jagged1 DNA Copy Number Variation Is Associated with Poor Outcome in Liver Cancer

Kazunori Kawaguchi, Masao Honda, Taro Yamashita, Hikari Okada, Takayoshi Shirasaki, Masashi Nishikawa, Kouki Nio, Kuniaki Arai, Yoshio Sakai, Tatsuya Yamashita, Eishiro Mizukoshi, and Shuichi Kaneko

From the Department of Gastroenterology, Kanazawa University Graduate School of Medical Science, Kanazawa, Japan

Accepted for publication
April 9, 2016.

Address correspondence to
Shuichi Kaneko, M.D., Ph.D.,
Department of Gastroenter-
ology, Kanazawa University
Graduate School of Medical
Science, 13-1 Takara-Machi,
Kanazawa, Ishikawa 920-8641,
Japan. E-mail: skaneko@m-kanazawa.jp.

Notch signaling abnormalities are reported to be involved in the acceleration of malignancy in solid tumors and stem cell formation or regeneration in various organs. We analyzed specific genes for DNA copy number variations in liver cancer cells and investigated whether these factors relate to clinical outcome. Chromosome 20p, which includes the ligand for Notch pathways, Jagged1, was found to be amplified in several types of hepatoma cells, and its mRNA was up-regulated according to α -fetoprotein gene expression levels. Notch inhibition using Jagged1 shRNA and γ -secretase inhibitors produced significant suppression of cell growth in α -fetoprotein-producing cells with suppression of downstream genes. Using *in vivo* hepatoma models, the administration of γ -secretase inhibitors resulted in reduced tumor sizes and effective Notch inhibition with widespread apoptosis and necrosis of viable tumor cells. The γ -secretase inhibitors suppressed cell growth of the epithelial cell adhesion molecule-positive fraction in hepatoma cells, indicating that Notch inhibitors could suppress the stem cell features of liver cancer cells. Even in clinical liver cancer samples, the expression of α -fetoprotein and Jagged1 showed significant correlation, and amplification of the copy number of Jagged1 was associated with Jagged1 mRNA expression and poor survival after liver cancer surgical resection. In conclusion, amplification of Jagged1 contributed to mRNA expression that activates the Jagged1-Notch signaling pathway in liver cancer and led to poor outcome. (*Am J Pathol* 2016, 186: 2055–2067; <http://dx.doi.org/10.1016/j.ajpath.2016.04.011>)

Globally, liver cancer has one of the poorest prognoses of all cancers.¹ Many liver cancer patients experience recurrence after effective treatments, including surgical resection at stage I, leading to high mortality rates. The causes of liver cancer include hepatitis B virus (HBV) and hepatitis C virus (HCV) infection, any other chronic liver injury such as primary biliary cirrhosis or autoimmune hepatitis, metabolic diseases such as nonalcoholic steatohepatitis or diabetes, and exposure to aflatoxin, a harmful chemical substance. Although the direct oncogenic factor of HCV-related liver cancer has not yet been discovered, HBV is reported to be associated with oncogenic factors of viral infection, including HBV-x genome. Moreover, a recently developed effective antiviral therapy for both HCV and HBV, using direct-acting agents in HCV or nucleos(t)ide analogues in HBV, has led to some patients developing liver cancer without developing advanced liver disease such as cirrhosis. However, this oncogenic mechanism has not been resolved

completely even in HBV and could become a major problem in the near future. Serologic tests are performed to diagnose liver cancer, to determine the appropriate therapies, and to monitor their effects. The most important serologic tumor markers are α -fetoprotein (AFP) or prothrombin induced by vitamin K absence-II. AFP is also associated with angiogenesis and with tumor cell growth, invasion, and metastasis. From hierarchical expression analysis using microarray, ephrin family genes have been found to be linked to AFP elevation or angiogenesis.² Serum AFP elevation or overexpression in the tissue can be seen in liver cancer samples; however, other proteins have also been discovered. For example, glypican 3 is significantly elevated and epithelial cell adhesion molecule (EpCAM) plays an

Supported by Viral Hepatitis Research Foundation of Japan (K.K.).

Disclosures: MACArray Karyo 1440 BAC-chip was supplied by Macrogen, Seoul, South Korea to K.K.

important role in liver cancer progenitor cells.^{3–5} Recent bioinformatics techniques have been applied to find proteins related to abnormal cellular signaling in liver cancer samples. We examined gene expression patterns using RNA samples of liver cancer tissue, including clinical samples, in addition to hepatoma cell lines or hepatoma cells in experimental small animals. At the DNA level, genomic alterations such as single nucleotide polymorphisms, loss of heterogeneity, or copy number amplifications or deletions were found, and these resulted in specific changes in liver cancer samples.^{6,7} Southern blot analysis, fluorescence *in situ* hybridization, and comparative genomic hybridization (CGH) have been used in cytogenetic studies. Recent progress in bioinformatics techniques has led to the discovery of specific genomic alterations in solid tumors such as tumors of the breast, colon, and pancreas, and these have been useful for diagnostics and in the selection of therapy. The same techniques have also resulted in the identification of specific genomic alterations in liver cancer. Altered genomic patterns in liver cancer influence patient mortality. A specific genomic location altered in liver cancer samples was reported to be associated with carcinogenesis or mitosis of the tumor cells. Previously, we reported that genomic alteration patterns affect the expression patterns in hepatoma cell lines and are associated with AFP production.⁸ Referring to related genes on microarray, we could estimate the types of genes involved and the candidate cell-cell interaction genes. Because it was important to determine whether a specific gene could be involved with a possible molecular therapeutic target or one of the pathways in a molecular target in clinical samples, we used microarray results to detect genomic abnormalities.

Materials and Methods

Hepatoma and Nonhepatoma Cell Lines

For genomic and gene expression analysis, we selected seven hepatoma cell lines: five cell lines, Huh7, HepG2, Hep3B, Huh6, and PLC/PRF/5, that produce AFP, and two cell lines, HLE and SKHep1, that do not. Human cervical cancer cell line HeLa and bile duct cancer cell line KMBC, neither of which produce AFP, were also analyzed as non-hepatoma cells. THLE-5b cells, a normal hepatocyte immortalized by using Simian virus 40, was selected for normal gene expression control for real-time detection PCR gene expression analysis.⁹

Clinical Liver Samples (Frozen Liver Cancer Samples)

Genetic analysis of frozen tumor samples from 38 liver cancer patients was conducted using real-time detection PCR. Most samples were collected during surgery. The patients included 17 who had HBV-related liver cancer and 18 who had HCV-related liver cancer; the remainder had other causes. We observed AFP expression in hepatoma

tissue samples, and it was up-regulated in approximately 37% of samples (six HBV, six HCV, two non-B and non-C). Surgery was performed on patients diagnosed from October 2000 to September 2006 at Kanazawa University Hospital. All patients gave informed consent for their tissues to be used for genomic analysis. By isolating genomic DNA from eight patients with elevated AFP expression (one sample was from an autopsy liver), we observed genomic copy number changes using real-time detection PCR by comparing the samples with noncancerous tissues.

Clinical Liver Samples

Formalin-fixed, paraffin-embedded samples from 115 liver cancer patients were selected, and genomic DNA was isolated by checking the tumor area under a microscope using similar hematoxylin and eosin–stained slices and scraping off tumor cells from samples on the glass slide. Genomic DNA isolation was performed using Min-elute formalin-fixed, paraffin-embedded kit (Qiagen, Venlo, the Netherlands). Surgery had been performed in patients diagnosed from 2004 to 2010 who gave consent for the genetic analysis of tissues. Of these samples, 31 were HBV-related liver cancer, 54 were HCV-related, one was both HBV and HCV infected, and 29 were non-HBV and non-HCV.

BAC Array Hybridization and Comparison with In-House cDNA Chip

Seven hepatoma cell lines and two nonhepatoma cell lines were used for bacterial artificial chromosome (BAC) microarray CGH. Manufactured microarray slides contained 1440 BAC clones, which covered most major chromosome regions (Macrogen, Seoul, South Korea). Genomic DNA (0.5 µg) from tumor cells was labeled by cyanine 5-dUTP (GE Healthcare Life Sciences, Uppsala, Sweden), and the same amount of control genomic DNA from peripheral blood mononuclear cells of healthy volunteers was labeled by cyanine 3-dUTP (GE Healthcare) and hybridized onto slides at 37°C overnight. The microarray technique was similar to the conventional method. A microarray scanner was used to calculate the intensity of each dye, and we analyzed the BAC clone spots in each chromosomal region using MAC Viewer (Macrogen) and viewed the log ratio chart. We compared the expression profiles from the results of the cDNA in-house microarray (Kanazawa chip version 2.1, 9600 spots) and analyzed the same cell line samples.⁸ These BAC array CGH data sets have been deposited in National Center for Biotechnology Information Gene Expression Omnibus (<http://www.ncbi.nlm.nih.gov/geo>; accession number GSE77224).

Real-Time Detection PCR of Notch-Related Genes

Real-time detection PCR was performed using cDNA from clinical samples or cell lines to measure the expression of Jagged1 and AFP. We also checked the expression levels of

other Notch-related genes, *Notch1* and *HES1*. We measured the genomic copy number variation (CNV) of Jagged1 using real-time detection PCR for genomic DNA samples. We used master mix reagent for real-time detection PCR (Life Technologies, Carlsbad, CA) and performed amplification using ABI7900 (Life Technologies). For CNV analysis, we used two different areas of the Jagged1 probe, and CNV calculation was performed using CopyCaller software version 2.0 (Life Technologies).

Cell Growth Assay after Jagged1 shRNA Silencing

We measured cell line growth after Jagged1 expression inhibition using shRNA vector (MISSION shRNA; Sigma-Aldrich, St. Louis, MO). Cell growth for up to 120 hours in Huh7, HepG2, HLE, and SKHep1 cells was compared using Jagged1 and a nontargeting negative control after infection by these lentivirus-transfected vectors. After puromycin selection, we compared cell growth using Cell Counting Kit-8 (Dojindo Laboratories, Kumamoto, Japan).

Notch Inhibition by Notch Inhibitors (GSIs)

To disrupt Notch signaling in hepatoma cells, we used two Notch inhibitors, L-685,485 and DAPT [(3,5-Difluorophenylacetyl)-Ala-Phg-OBu^t; Peptide Institute, Inc., Minou, Japan]. These drugs, γ -secretase inhibitors (GSIs), are administered to prevent amyloid β accumulation and have been used in clinical trials in Alzheimer patients. In addition to Alzheimer research, many trials are ongoing for solid malignant tumors, such as breast cancers, lung cancer, and colorectal cancers.^{10,11} L-685,485 strongly inhibits amyloid β ; thus, it is widely used in basic research, whereas, despite similar inhibitory effects, DAPT needs a greater concentration to inhibit signal peptide peptidase.^{12,13} We observed cell proliferation in four hepatoma cell lines (Huh7, HepG2, HLE, and SKHep1) after administering 10 μ mol/L of these drugs. We measured data from five wells for each sample and calculated absorption to measure cellular growth. We changed the medium at 24, 48, 72, 96, and 120 hours and checked cell growth at 0, 72, 120, and 168 hours (Huh7 and SKHep1) or 0, 48, 96, and 168 hours (HepG2 and HLE). The reason for varying the measurement times was that cellular growth differed, depending on cell lines, and we could not estimate the confluent state for these cell sets. Moreover, we checked the expression of the downstream gene, *HES1*, in Huh7 and HLE by real-time detection PCR.

Cell Separation of EpCAM⁺ and EpCAM⁻ Huh7 Cells

Because Huh7 cells include the heterogenic EpCAM⁺ stem cell feature and this stem cell population is associated with the malignant characteristics of cells, we separated EpCAM⁺ and EpCAM⁻ fractions using fluorescence-activated cell sorting (FACS Aria III; BD Franklin Lakes, NJ). Cells were spread onto culture dishes after sorting and

incubated at 37°C in 5% CO₂. After several passages, we administered 10 μ mol/L L-685,485 and DAPT. Relative cellular growth rates were analyzed using Cell Counting Kit-8 (Dojindo Laboratories).

GSIs Treatment for NOD-SCID Mouse Hepatoma Models

Seven-week-old nonobese diabetic severe combined immunodeficiency (NOD-SCID) mice (NOD.CB17-Prkdc<SCID>/J, male; Charles River Laboratories, Wilmington, MA) were given subcutaneous injections of 1×10^5 of Huh7 cells. Percutaneous administration of 5-mg/kg doses of L-685,485 and 20-mg/kg doses of DAPT dissolved in dimethyl sulfoxide (DMSO) or DMSO alone (control) was started. To compare tumor size, we injected these drugs three times per week, sacrificed the mice after 2 weeks, and measured the hepatomas. To assess survival, we injected drugs twice a week and observed the mice for up to 6 weeks after the injections. Thereafter, we sacrificed the remaining live mice and applied immunofluorescence. Several specific polyclonal antibodies were added to paraffin-embedded tumor samples before staining with 3,3'-diaminobenzidine tetra hydrochloride (DAB) and hematoxylin. We calculated the CK19-stained areas in the cells and compared the control and GSI-treated mouse tumor samples using ImageJ software version 1.47v (NIH, Bethesda, MD; <https://imagej.nih.gov/ij>).

Results

20p Genomic Amplification of Hepatoma Cell Lines Using Array-Based CGH

We analyzed the genomic amplification and deletion of hepatoma cells using BAC microarray CGH. Five of the cell types produce AFP (Huh7, Hep3B, Huh6, PLC/PRF/5, HepG2), whereas two do not (SKHep1 and HLE). The genomic DNA of these cells was previously analyzed using cDNA microarray CGH, and it was found that AFP production is associated with a different genomic cluster of cells.⁸ A similar result was observed in cDNA and BAC array CGH, but some results were different. This was because the target DNA on the array slide was different and exons but not introns were present, in contrast to genomic DNA; thus, the intron sequence was skipped. Moreover, the length of the cDNA was shorter than the BAC clone. The genomic DNA was amplified sequentially in some chromosome locations and deleted in others. The 20p position was particularly amplified compared with other chromosome locations, although we found other genomic alterations at other locations (Figure 1A). Because we have already reported that AFP-positive and -negative hepatoma cells can differentiate between genomic amplification and expression, we mixed the data of each spot from the BAC array and calculated the average AFP-positive and -negative characteristics (Figure 1B). The copy number differed at other chromosomal locations based on AFP-producing

status—these differences were strong, especially in the 20p location.

Amplification and Overexpression of Jagged1 Gene in AFP-Positive Hepatoma Cells

We selected genes in each of the BAC clones to focus on 20p in the BAC microarray (Table 1). Although not all spots in the BAC array expressed genes, several were found

during the analysis. From 9800 in-house genes of cDNA microarray, several were located in the 20p area, and we compared the average expression ratio in AFP-positive and -negative cell lines for some genes. We found that the Jagged1 gene showed a distinct pattern (Table 1, Figure 1A). The underlined genes in Table 1 are listed on the in-house 9800 microarray. The expression of other genes was not up-regulated in AFP-positive cells. With the use of cDNA microarray, we previously showed that

Figure 1 *In vitro* genomic copy number and expression analysis in hepatoma or nonhepatoma cells. **A:** Genomic copy number changes of chromosome 20 in each cell line using BAC microarray. Copy number changes in each chromosome area were calculated using commercial software. Human hepatoma cell lines (Huh7, HepG2, Huh6, Hep3B, PLC/PRF/5, HLE, and SKHep1) and human nonhepatoma cell lines (KMBC and HeLa) were analyzed by comparing with healthy human volunteer PBMCs as control. Red dotted lines with arrows indicate Jagged1 gene copy number changes in 20p12.2. Values above zero show copy number gains and those below indicate copy number losses. Average log2 (cyanine 3/cyanine 5) signal ratios of BAC clones were calculated for each sample, and a threshold of 0.25 log2 ratio units was used for defining copy number gains and losses. **B:** Mean copy number changes between AFP-positive (yellow) and -negative cells (green). **C:** Real-time detection PCR analysis of Jagged1 gene in cell lines. These values are log-based Jagged1/ β -actin expression ratios. Huh7, Hep3B, HepG2, PLC/PRF/5, and Huh6 cells are AFP producing, whereas SKHep1, HLE, and HeLa are not. **D:** Real-time detection PCR analysis of Jagged1 genome in cell lines. These values were calculated by comparing changes with genomic changes in healthy PBMCs. BAC, bacterial artificial chromosome; PBMC, peripheral blood mononuclear cell.

Table 1 BAC Clones of Chromosome No. 20 on Microarray Slides and Representative Genes

BAC clone ID	Chromosome location	BAC size	Representative genes	GenBank ID*	Average expression ratio of AFP+ cells/ AFP- cells
BAC40_E05	20p13	110124	<i>LOC343637, RSP04</i>		
BAC109_E03	20p13	97397	<i>ATRN, GFRA4, ADAM33</i>	NM_139321	0.88
BAC127_K22	20p12.3	91214	<i>C20orf30, PCNA, CDS2</i>	NM_002592	0.98
BAC85_D15	20p12.3	100000			
BAC155_K05	20p12.3	90398	<i>PLCB1</i>		
BAC143_E21	20p12.2	98802	<i>PLCB4</i>	NM_000933	1.08
BAC86_O12	20p12.2	136522	<i>JAG1</i>	NM_000214	2.14
BAC165_M10	20p12.1	88468			
BAC86_O11	20p12.1	153399	<i>MACROD2, CR596518</i>		
BAC81_M07	20p12.1	126920	<i>MACROD2</i>		
BAC81_K08	20p12.1	88837	<i>KIF16B</i>		
BAC29_A11	20p12.1	131740	<i>RPL7AL3, SNRPB2, OTOR</i>	NM_003092	0.4
BAC132_A05	20p12.1	93613			
BAC96_K10	20p12.1	112559	<i>PCSK2, BFSP1</i>	NM_002594	0.99
				NM_001195	0.98
BAC44_G02	20p11.23	110517	<i>DTD1</i>		
BAC75_H16	20p11.23	108696	<i>KIAA1272, RALGA1A2 C20orf74, AK026194</i>		
BAC63_I12	20p11.22	104597	<i>NKX2-2</i>		
BAC110_E03	20p11.21	113928	<i>VSX1, ACSS1</i>		
BAC36_M13	20p11.21	173706	<i>ENTPD6, PYGB, C20orf22, PPIAP2, BC128043, ABHD12, CD39L2</i>	NM_001247	1.6
			<i>CBFA2T2</i>	NM_002862	0.79
BAC204_A05	20q11.21	89320			
BAC140_D09	20q11.23	125023			
BAC43_F22	20q12	171393	<i>LOC339568</i>		
BAC239_E02	20q12	91366	<i>CR612573, CR593014</i>		
BAC97_K09	20q12	76966	<i>LPIN3, EMILIN3, KIAA1335, CH06</i>		
BAC21_G16	20q12	109908			
BAC52_H17	20q12	114526	<i>PTPRT</i>	NM_133170	0.87
BAC75_I24	20q13.12	86258	<i>ADA, AK090842, CR597563</i>	NM_000022	0.71
			<i>CR616794, WISP2</i>	NM_003881	1.11
BAC12_K18	20q13.13	109372	<i>STAU1</i>		
BAC193_H13	20q13.13	102377			
BAC72_N10	20q13.2	95499			
BAC51_F06	20q13.2	224038			
BAC97_O05	20q13.2	102916			
BAC63_G09	20q13.32	121623	<i>PPP4R1L, RAB22A</i>		
BAC142_K22	20q13.32	81986	<i>PHACTR3, SYCP2</i>		
BAC42_C03	20q13.33	102953	<i>CDH4, AK097866</i>	NM_001794	1.11

Underlined representative genes; 9800 in-house cDNA microarray listed genes.

*GenBank (<http://www.ncbi.nlm.nih.gov/genbank>).

AFP, α -fetoprotein; BAC, bacterial artificial chromosome.

AFP-positive and -negative cells had differential gene expression and genomic alterations.⁵ We focused on the Jagged1 gene and analyzed its role in hepatoma cells or clinical samples. With the use of the real-time detection PCR method, Jagged1 genes showed distinct up-regulation in AFP-positive cell lines and down-regulation in AFP-negative cell lines (Figure 1C). Furthermore, the data showed that Jagged1 gene was not up-regulated in non-hepatoma cell lines such as HeLa. We analyzed alterations in the Jagged1 copy number using real-time detection PCR and found that most of the cell lines showed similar expression, except HLE cells (Figure 1D).

Cell Growth Suppression in AFP-Producing Hepatoma Cells after Jagged1 shRNA Silencing

Strong suppression of cell growth in Huh7 and HepG2 cell lines was found after shRNA silencing of Jagged1 gene (*Jag1*) (Figure 2). However, AFP-negative cells, HLE and SKHep1, showed no growth suppression. We also performed Jagged1 siRNA knockdown and evaluated cell growth; however, no significant suppression was observed, even in Huh7 and HepG2 cells (data not shown). Real-time detection PCR confirmed that downstream *HES1* gene expression was suppressed in Huh7 for Jagged1 siRNA

Figure 2 Jagged1 inhibition of hepatoma cells impacts on tumor growth *in vitro*. Cell growth after Jagged1 shRNA transfection in Huh7, HepG2, HLE, and SKHep1 cells. Huh7 and HepG2 are α -fetoprotein-producing cells, and HLE and SKHep1 are nonproducing cells. In Jagged1 shRNA transfected cells (red), cellular growth was compared with control shRNA-transfected cells (black). *** $P < 0.001$.

(Supplemental Figure S1). The reason for this result may be because the induction of Jagged1 knockdown by siRNA was not efficient and successful transfection might only have occurred in a small number of cells.

Effect of Notch Signal Inhibition of Hepatoma Cells by GSIs

Two types of GSIs, L-685,458 and DAPT, were used for the cellular reaction of Notch pathway inhibition among hepatoma cell lines. L-685,458 and DAPT diluted with DMSO were added to cells, and cell proliferation was compared with the control (DMSO alone). Inhibition occurred in Huh7 and HepG2, which are AFP-producing cell lines, whereas there was no significant inhibition in non-AFP-producing cell lines, HLE and SKHep1 (Figure 3A). We calculated concentration that inhibits 50% (IC_{50}) values for these cells, and the values of L-685,458-treated groups were smaller than those treated with DAPT and were smaller in Huh7 cells than in other cells (Table 2). Notch-related gene expression after 120 hours showed that HES1 expression was suppressed using the two types of GSIs in both Huh7 and HLE (Figure 3B). However, other Notch signaling-related genes, Jagged1, *Notch1*, and *Notch2*, were not altered apart from *Notch1* expression after DAPT treatment in HLE, suggesting that only downstream genes are affected by GSIs.

Effectiveness of GSIs in EpCAM⁺ Cancer Stem Cells

Notch signaling plays a role in the functions of stem cells, and EpCAM⁺ is strongly associated with cancer stem cells in hepatomas. Cell growth was compared after GSIs treatment in EpCAM⁺ and EpCAM⁻ Huh7 cell fractions after separation using a cell sorter. We observed significant cell growth suppression in EpCAM⁺ cells with DAPT, and L-685,458

suppressed both EpCAM⁺ and EpCAM⁻ features, suggesting that GSIs have antitumor effects, even in hepatoma cells with the EpCAM⁺ cancer stem cell feature (Figure 3C).

Antitumor Effect of Notch Inhibitors in Mouse Hepatoma Models

Slower tumor formation was observed after 2-week administration of Notch inhibitors, L-685,458 and DAPT (Figure 4A). The effect was more pronounced for DAPT ($P = 0.04$), and, after a longer period of observation, there was a tendency for earlier death in control cases (Figure 4B). The dynamic tumor status in GSIs cases after longer administration showed tumor necrosis and apoptosis (Figure 4C). We found that EpCAM staining was weak in the GSIs groups except in necrotic areas, suggesting GSIs had inhibited EpCAM production (Figure 4C). Moreover, CK19 staining was significantly reduced in GSIs cases ($P < 0.001$), which indicated that the malignant characteristics of the hepatoma cells were controlled (Figure 4D). Brighter hematoxylin staining in GSIs-treated cases indicated that HES1 staining was diminished in the nucleus (Figure 4C). Jagged1 and Notch1 were stained by immunofluorescence in control tumor samples; however, these stains were not changed in the L-685,458 or DAPT treatment groups. This result showed that ligands and receptors were not affected by GSIs (Supplemental Figure S2A). This is reasonable data; we checked real-time detection PCR data of HES1 and found a decrease after GSIs treatment (Supplemental Figure S2B).

Association between Amplification of Jagged1 Genome in Liver Cancer Tissues and Poor Prognosis

Genomic DNA in formalin-fixed, paraffin-embedded samples from surgically resected liver cancer cases was analyzed, and

Figure 3 Notch inhibitor impact on AFP-producing and EpCAM⁺ hepatoma cells *in vitro*. **A:** Relative cellular growth curve of hepatoma cell lines after administration of γ -secretase inhibitors L-685,458 and DAPT. We administered 10 μ mol/L of each reagent dissolved in DMSO at 0 hour, 72 hours, 120 hours, and 168 hours and blank reagents that included only DMSO (control). **B:** Relative expression of HES1, Jagged1, Notch1, and Notch2 after L-685,458 and DAPT treatment in Huh7 and HLE cells. Total RNA was isolated after 120 hours administration. **C:** Relative cellular growth of L-685,458- or DAPT-treated Huh7 cells after separation of EpCAM⁺ and EpCAM⁻ fractions using fluorescence-activated cell sorting. We compared 120 hours of cellular growth in each with the DMSO control. Data are expressed as means $\pm$ SD. * $P < 0.05$; *** $P < 0.001$ control versus L-685,458 and DAPT. AFP, α -fetoprotein; DAPT, (3,5-Difluorophenylacetyl)-Ala-Phg-OBu^t; DMSO, dimethyl sulfoxide; EpCAM, epithelial cell adhesion molecule.

the cases were followed for up to 5 years after surgery. High Jagged1 CNV cases (≥ 1.5 -fold) and low CNV cases (< 1.5 -fold) were divided and the clinical characteristics were compared. Advanced TMN classification and positive vascular invasion were associated with high Jagged1 CNV (Table 3). Multivariate analysis showed that high Jagged1 CNV (≥ 1.5 -fold) in cases of liver cancer recurrence after surgical resection was associated with survival (Table 4). Copy number alterations were analyzed using two different areas of the Jagged1 genome locus, and the proximal copy number multiple of 0.5 was calculated. CNV analysis of the

Table 2 IC₅₀ Values of L-685,458, and DAPT in Hepatoma Cell Lines

Cell line	IC ₅₀ (μ mol/L)	
	L-685,458	DAPT
Huh7	12.91 $\pm$ 0.55	118.26 $\pm$ 40.42
HepG2	12.69 $\pm$ 4.60	155.27 $\pm$ 7.58
HLE	21.76 $\pm$ 0.84	244.86 $\pm$ 7.95
SKHep1	12.18 $\pm$ 2.90	153.24 $\pm$ 14.38

DAPT, (3,5-Difluorophenylacetyl)-Ala-Phg-OBu^t; IC₅₀, concentration that inhibits 50%.

Jagged1 locus in 110 liver cancer samples revealed that high CNV cases (≥ 1.5 -fold, 65 cases) had significantly lower survival rates ($P = 0.019$) compared with low CNV cases (< 1.5 -fold, 45 cases) (Figure 5A). We also classified three groups (CNV ≥ 2.0 -fold, 28 cases; $2.0 \geq$ CNV > 1.5 , 37 cases; CNV < 1.5 -fold, 45 cases), and higher CNV groups showed poorer survival rates compared with lower CNV cases ($P = 0.014$) (Figure 5B). AFP elevation status was analyzed because Jagged1 amplification is observed especially in AFP-producing hepatoma cells *in vitro*. We classified the two groups according to the presence (≥ 100 ng/mL) or absence of AFP elevation. There was significantly higher Jagged1 CNV in AFP-elevated cases (Figure 5C).

The expression of AFP and Jagged1 genes among frozen resected liver cancer tissues were compared using real-time detection PCR. AFP and Jagged1 expression showed significant correlation after the selection of AFP-up-regulated samples ($P < 0.05$). Moreover, there was a strong correlation between AFP expression and Jagged1 genome amplification ($P < 0.01$) (Figure 5, D and E). Jagged1 expression and Jagged1 genome amplification also correlated ($P < 0.05$) (Figure 5F), suggesting that the AFP value was

Figure 4 Antitumor effect of γ -secretase inhibitors in nonobese diabetic severe combined immunodeficiency mouse models implanted with Huh7 cells. **A:** Tumor size comparison between L-685,458, DAPT, and negative control after 2 weeks of treatment. Tumor diameters of each of the five tumor samples were analyzed. **B:** Cumulative survival rate compared between GSIs (L-685,458 and DAPT) and control cases. Survival rates of each of the five samples were compared and calculated using the log-rank t -test. **C:** Immunostaining using anti-caspase 3, CK19, HES1, and EpCAM antibodies for control and L-685,458-treated tissues. Immunostaining was performed using diaminobenzidine, and cellular nuclei were stained by hematoxylin. **D:** Quantification of CK19⁺ between control and GSIs. Calculations were performed for each of five CK19-immunostained slides. * $P < 0.05$, ** $P < 0.01$, and *** $P < 0.001$. Scale bar = 100 μ m. DAPT, (3,5-Difluorophenylacetyl)-Ala-Phe-OBu^t; EpCAM, epithelial cell adhesion molecule; GSI, γ -secretase inhibitor.

associated with Jagged1 expression and genome amplification in clinical samples. We also confirmed a correlation between Jagged1 and Notch1 expression by real-time detection PCR using these samples. We assessed Notch1 and downstream HES1 genes and found a strong correlation, which suggested that Notch signaling was up-regulated in AFP-positive clinical tissues (Supplemental Figure S3).

Discussion

Genomic alterations such as genome copy number gain or loss in cancerous cells can be analyzed using microarray

based-CGH.^{6,14} Amplified or deleted regions are found in various types of cancers, but these specific abnormal regions differ in different organs.¹⁵ Some reports show that regions of genomic alterations in cancer are associated with expression abnormalities.^{16,17} We previously reported a significant association between altered genomic copy number and gene expression patterns in hepatoma cells and, using hierarchical clustering analysis by microarray, found that the factor involved was AFP-producing status.⁸ Chromosome 20p locations in hepatoma cells apparently differed in terms of genomic copy number, and AFP-producing hepatoma cells differed from non-AFP-producing cells.

Table 3 Clinical Characteristics of Liver Cancer Patients Who Had Undergone Surgical Resection

Clinical category	CNV ≥ 1.5	CNV < 1.5	P
No. of patients	68	47	
Sex, male vs female	50 vs 18	39 vs 8	0.234
Age, years	63 (38–82)	66 (44–80)	0.054
Viruses, HBV/HCV/non-B non-C	21/33/15*	11/22/14	0.540
Child Pugh score, 5/6/ ≥ 7	55/5/6 [†]	36/7/1 [‡]	0.054
AFP, ≥ 100 ng/mL vs < 100	27/41	11/36	0.067
PIVKA-II, (≥ 100 mAU/mL vs < 100)	35/31	26/19	0.622
Histologic grading, moderate to poor vs well to moderate	57/11	37/7	0.970
Tumor size, mm	45 (7–230)	35 (10–150)	0.359
Tumor morphology, multiple vs uni	26/42	13/34	0.239
Vascular invasion, positive vs negative	15 vs 53	1 vs 46	0.002**
Liver cancer recurrence, positive vs negative	31 vs 26	32 vs 12	0.059

Data are expressed as *n* or median (range) unless otherwise indicated.

*One patient had a double infection of HBV and HCV.

***P* < 0.01.

[†]The score for two patients could not be calculated.

[‡]The score for three patients could not be calculated.

AFP, α -fetoprotein; CNV, copy number variation; HBV, hepatitis B virus; HCV, hepatitis C virus; PIVKA-II, prothrombin induced by vitamin K absence-II; uni, unifocal.

A previous report showed genomic alterations in liver cancer using CGH analyses and frequent gains were observed in 1q, 6p, 8q, and 17q, whereas losses were observed in 4q, 8p, 13q, 16q, and 17p.¹⁸ In addition to these gained locations, 20p11 was also present in some liver cancer tissue.¹⁹ Moreover, five BAC clones for this experiment were found listed in BAC array CGH, but only Jagged1 gene was a candidate because the different expression patterns were associated with the AFP-producing status of hepatoma cells using microarray.

Notch signaling pathway is widespread and essential for cell-cell interactions or intracellular transduction by Notch intracellular domain.²⁰ Notch ligands and receptors perform major roles in this pathway, and abnormalities such as point mutations are reported to be the cause of Alagille syndrome, a systemic disorder in humans.^{21,22} In the oncology field, Notch signaling abnormalities are related to carcinogenesis or tumor vessel abnormalities.^{23,24} Notch signaling-targeted therapy has been developed in the field of neural diseases such as Alzheimer disease. One of the Notch inhibitors, GSI, restricts cleavage of the Notch receptor in the cell membrane and disturbs signal transduction by Notch intracellular domain. Clinical trials were under way, but reports of gastrointestinal toxicity halted further study.²⁵ This drug is useful for inhibiting aggregation of amyloid protein in Alzheimer disease,²⁶ although its effectiveness has been reported in the oncology field in targeted molecular therapy.²⁷ For example, hematopoietic tumor therapy is highly effective, and various clinical studies in other solid tumors are ongoing.^{28–30}

Some genomic alteration studies have already been performed in liver cancer samples, and many reports using array CGH method or other new bioinformatics techniques have been published.³¹ Genome amplification of cancer cells usually shows that the check point system for DNA repair was destroyed, leading to gene amplification, especially in malignant cells, including those of liver cancer. However, it is important to consider that liver cancer includes several histologic types and genomic alteration patterns differ by type; variations occur in differentiation, stem cell features, AFP production, and so forth. The causes also differ because advanced chronic liver injury can be caused by HBV, HCV, or other factors. Some reports suggest that genomic abnormalities in liver cancer cause poor survival and candidate genes have been reported; however, they are not associated with the development of any clinical drugs.³² Our method selected genes by microarray analysis, but we concluded that these candidate genes would be important for future effective liver cancer therapy. The relation between genomic and

Table 4 Cox Regression Analysis of Clinical and Genomic Variables Associated with Survival in Liver Cancer

Variable	Univariate	Multivariate	
	P	P	HR (95% CI)
Sex, male vs female	0.824	0.104	2.639 (0.820–8.497)
Age, ≥ 60 years vs < 60 years	0.551	0.612	1.258 (0.519–3.050)
Jagged1 CNV, ≥ 1.5 vs < 1.5	0.002**	0.018*	3.494 (1.243–9.816)
AFP, ≥ 100 vs < 100	0.079	0.187	1.897 (0.733–4.909)
PIVKA II, ≥ 100 vs < 100	0.001**	0.930	0.956 (0.353–2.591)
Liver cancer recurrence, yes vs no	0.139	0.001**	9.032 (2.472–33.00)
Child-Pugh scores, ≥ 7 vs < 7	0.024*	0.113	2.704 (0.789–9.268)
Histologic grading, moderate to poor vs well to moderate	0.221	0.371	2.000 (0.438–9.146)
Tumor size, ≥ 30 mm vs < 30 mm	0.003**	0.060	3.073 (0.954–9.899)
Vascular invasion, positive vs negative	0.001**	0.553	1.423 (0.443–4.574)

P* < 0.05, *P* < 0.01.

AFP, α -fetoprotein; CNV, copy number variation; HR, hazard ratio; PIVKA-II, prothrombin induced by vitamin K absence-II.

Figure 5 Amplified Jagged1 liver cancer patients have poor survival. We analyzed Jagged1 CNV in liver cancer tissue in formalin-fixed, paraffin-embedded and frozen tissue to compare genomic alterations and expression using real-time detection PCR. **A:** Overall survival data after surgical resection of liver cancer were divided into two groups (CNV < 1.5 and ≥ 1.5). The P value was calculated using log-rank t -test. We followed patients for up to 60 months after surgery. **B:** Overall survival data divided into three groups (CNV < 1.5 , CNV = 1.5, and CNV ≥ 2). **C:** Comparison of CNV values in two groups of high (≥ 100 ng/mL) and low (< 100 ng/mL) AFP using the U test. We used the day of surgery as the approximate date that the serum AFP values were obtained. **D:** Relation between Jagged1 and AFP expression by real-time detection PCR using frozen liver cancer tissues. We analyzed only AFP up-regulated liver cancer tissues. **E:** Relation between Jagged1 genomic alterations and AFP expression. **F:** Relation between Jagged1 genomic alterations and expression. * $P < 0.05$, ** $P < 0.01$. ACTB, β -actin; AFP, α -fetoprotein; CNV, copy number variation.

expression alterations can explain their oncologic significance. We investigated the importance of AFP-producing status in hepatoma cells using genome microarray analysis of expression. We focused on Jagged1 gene because it was both amplified and overexpressed in AFP-positive hepatoma cells. In another report, Jagged1 expression was associated with HBV-x genes³³; however, we included HCV-infected clinical liver cancer cells. Mutations in Jagged1 gene are reported to cause abnormalities of the bile duct cells and Alagille syndrome in the gastroenterology field. However, an association with cancer has not been investigated. By conducting Notch-related basic research, Notch ligand and receptor expression abnormalities have been linked to several types of cancers. Jagged1 gene expression abnormalities in liver cancer have already been reported, although copy number changes have not been discussed.³⁴ However, recent cancer research has revealed that copy number changes in specific genes are associated with carcinogenesis or the aggressiveness of the cancer and correlate with patient survival.^{35,36} In the hepatologic field, some reports have mentioned links between genomic alterations, severity of liver cancer, and survival.^{37,38} Our study revealed a correlation between AFP elevation and Jagged1 genomic copy number changes. From array CGH data and PCR-based CNV assay, Jagged1 CNV was not reduced in HLE cells even though these are not AFP-producing cells. This means Notch signaling, including Jagged1 underexpression, is applicable to non-AFP-producing cells, but the decrease in

Jagged1 CNV does not fully correlate in these cells. These data indicated that HLE cells are an exception, but Notch transduction suppression was significant. Furthermore, the data showed that CNV value was not elevated, which indicated that Jagged1 CNV did not change in these cells, and underexpression of Jagged1 was not directly associated with a decrease in Jagged1 CNV. We analyzed the genomic alterations of Jagged1 genes and AFP elevation using surgically resected samples. AFP elevation and overexpression of Jagged1 gene were calculated and related to liver cancer cases with elevated AFP. With the use of surgical samples, real-time detection PCR showed that AFP expression and Jagged1 genomic copy number changes also correlated. We concluded that the genomic copy numbers of Jagged1 gene were associated with AFP expression. Notch-related gene expression analysis after Jagged1 knockdown indicated that Notch signal transduction affected *HES1* gene by modulation of the Notch ligand.

In hepatoma cells, GSIs can be used both *in vitro* and *in vivo*. The results showed that, compared with the control, these drugs significantly inhibited growth in AFP-producing hepatoma cells. However, growth was strongly inhibited using DAPT but not L-685,458 in KMBC cell lines (data not shown). This result suggested that any bile duct cell-related cause could be efficiently suppressed by the administration of Notch inhibitors.

We used two types of Notch inhibitor L-685,458 and DAPT. The former is a more efficient signal peptide

peptidase inhibitor than DAPT, and signal peptide peptidase is essential for HCV replication.¹² This inhibitor does not directly inhibit Jagged1 because it is a ligand in the cells and in cell interactions, but, from our results, there appears to be some association, which will be investigated in further *in vivo* experiments. Recent clinical oncology research showed that the chemical was more effective in combination with currently used drugs, such as sorafenib, for advanced stages of liver cancer than as a monotherapy.^{39–41} It may provide effective data for *in vitro* or *in vivo* models. Because Notch signaling exists even in non-AFP-producing cells, it can suppress downstream *HES1* genes by GSIs in, for example, HLE cells. Because GSIs catalyze the membrane protein that affects Notch1 cleavage, they may not reduce Jagged1, Notch2, or even Notch1. Hence, our data imply that the expression of upstream Jagged1 expression and other receptors are not affected. Furthermore, the reason why Notch1 was significantly suppressed by DAPT only in HLE might be because it was suppressed in both Huh7 and HLE cells, and showed Notch1 consumption by not restricting Jagged1-Notch1 signaling, especially in HLE. We demonstrated an antitumor effect in AFP-producing hepatoma cells and that GSIs caused cell growth suppression even in EpCAM⁺ Huh7 cells. These data suggest that GSIs could be aimed toward the more malignant features of cells as promising new targeted molecular therapy. Liver progenitor cells and cancer stem cells are sensitive to GSIs treatment based on our experiments. Previous studies show that hepatoma cell lines and liver cancer tissues are heterogeneous, and cancer stem cells expressing EpCAM can be separated.^{3,42} Our experiments showed that antitumor effects occurred in the EpCAM⁺ fraction *in vitro*, although it is difficult to observe precise reactions in clinical tissues. DAPT administration resulted in antitumor effects only in the EpCAM⁺ cell fraction. Because these cells are particularly malignant, we would expect good prognoses after GSIs administration.

We used NOD-SCID mouse models implanted with human hepatoma cells and performed GSIs treatment. Tumor formation was suppressed and survival was extended by treatment, which indicates that *in vivo* Notch suppression causes effective antitumor effects in hepatomas. Although metastasis was not observed, earlier death because of tumor enlargement and more dominant viable tumor cells occurred in control cases. GSIs were dissolved in DMSO, which was also used in control cases. Although DMSO shows cytotoxicity in animal models, the amount (50 μ L/animal) used in this study did not affect survival rates.⁴³

Jagged1 CNV analysis of surgically resected liver cancer patients showed that amplification was associated with poor survival, and copy number gain resulted in poor prognosis. Genomic copy number alterations are associated with survival, and this result indicated that these changes were associated with malignant characteristics and influenced patient survival. This region may exist in other regions of the liver cancer genome. Statistical analysis showed that

high AFP levels are associated with high Jagged1 CNV. Moreover, we found significantly shorter overall survival in higher Jagged1 CNV cases that were dependent on CNV values. A comparison of CNV ≥ 1.5 and <1.5 groups, showed that AFP values were unchanged; even if CNV was high at surgical resection AFP is not because it is an earlier stage. However, even if AFP is not elevated and Jagged1 CNV is high, the clinical course is worse, earlier recurrences of liver cancer occur, and AFP rapidly increases at the end stage. However, when AFP is high at surgical resection, many cases show high Jagged1 CNV; conversely, when AFP is low, although some patients have higher CNV, in many cases it is lower and the clinical course is quite good (Figure 5C).

We explained that Jagged1 genomic amplification and overexpression is related to Notch signal enhancement and AFP positivity and the malignant characteristics of liver cancer cells were related to these abnormalities. Although the mechanism of AFP production and Notch enhancement were not directly explained, Notch signal transduction was associated with the activation of cancer stem cells, including those in liver cancer, and we found that the EpCAM⁺ fraction in hepatoma cells are sensitive to Notch signal inhibition. Advanced and end-stage liver cancer clinical cases show much higher AFP levels and most show Notch activation, including Jagged1 and Notch1 overexpression. Notch pathways are related to angiogenesis, and AFP-up-regulated cases show higher levels of angiogenic factors, such as the Ephrin family reported previously by our group; thus, it is reasonable to expect Notch signaling activation in AFP-up-regulated cases.² Epidemiologic studies show that liver cancer has a poor prognosis; therefore, it is important to focus on molecules that are well-known liver tumor markers, such as AFP.

Our results showed that the Jagged1 genomic region and Notch activity are associated with the malignant characteristics of liver cancer and can predict the outcome of liver cancer therapy.

Acknowledgments

We thank Mikiko Nakamura, Masayo Baba, and Nami Nishiyama for technical assistance.

Supplemental Data

Supplemental material for this article can be found at <http://dx.doi.org/10.1016/j.ajpath.2016.04.011>.

References

1. Yang JD, Roberts LR: Hepatocellular carcinoma: a global view. *Nat Rev Gastroenterol Hepatol* 2010, 7:448–458
2. Iida H, Honda M, Kawai HF, Yamashita T, Shiota Y, Wang BC, Miao H, Kaneko S: Ephrin-A1 expression contributes to the malignant

- characteristics of {alpha}-fetoprotein producing hepatocellular carcinoma. *Gut* 2005, 54:843–851
3. Yamashita T, Ji J, Budha A, Forgues M, Yang W, Wang HY, Jia H, Ye Q, Qin LX, Wauthier E, Reid LM, Minato H, Honda M, Kaneko S, Tang ZY, Wang XW: EpCAM-positive hepatocellular carcinoma cells are tumor-initiating cells with stem/progenitor cell features. *Gastroenterology* 2009, 136:1012–1024
 4. Yamashita T, Honda M, Nio K, Nakamoto Y, Takamura H, Tani T, Zen Y, Kaneko S: Oncostatin m renders epithelial cell adhesion molecule-positive liver cancer stem cells sensitive to 5-Fluorouracil by inducing hepatocytic differentiation. *Cancer Res* 2010, 70:4687–4697
 5. Kawai HF, Kaneko S, Honda M, Shiota Y, Kobayashi K: Alpha-fetoprotein-producing hepatoma cell lines share common expression profiles of genes in various categories demonstrated by cDNA microarray analysis. *Hepatology* 2001, 33:676–691
 6. Pollack JR, Perou CM, Alizadeh AA, Eisen MB, Pergamenschikov A, Williams CF, Jeffrey SS, Botstein D, Brown PO: Genome-wide analysis of DNA copy-number changes using cDNA microarrays. *Nat Genet* 1999, 23:41–46
 7. Pollack JR, Sorlie T, Perou CM, Rees CA, Jeffrey SS, Lønning PE, Tibshirani R, Botstein D, Borresen-Dale AL, Brown PO: Microarray analysis reveals a major direct role of DNA copy number alteration in the transcriptional program of human breast tumors. *Proc Natl Acad Sci U S A* 2002, 99:12963–12968
 8. Kawaguchi K, Honda M, Yamashita T, Shiota Y, Kaneko S: Differential gene alteration among hepatoma cell lines demonstrated by cDNA microarray-based comparative genomic hybridization. *Biochem Biophys Res Commun* 2005, 329:370–380
 9. Pfeifer AM, Cole KE, Smoot DT, Weston A, Groopman JD, Shields PG, Vignaud JM, Juillerat M, Lipsky MM, Trump BF, Lechner JF, Harris CC: Simian virus 40 large tumor antigen-immortalized normal human liver epithelial cells express hepatocyte characteristics and metabolize chemical carcinogens. *Proc Natl Acad Sci U S A* 1993, 90:5123–5127
 10. Espinoza I, Miele L: Notch inhibitors for cancer treatment. *Pharmacol Ther* 2013, 139:95–110
 11. Takebe N, Nguyen D, Yang SX: Targeting notch signaling pathway in cancer: clinical development advances and challenges. *Pharmacol Ther* 2014, 141:140–149
 12. Li YM, Xu M, Lai MT, Huang Q, Castro JL, DiMuzio-Mower J, Harrison T, Lellis C, Nadin A, Neduvilil JG, Register RB, Sardana MK, Shearman MS, Smith AL, Shi XP, Yin KC, Shafer JA, Gardell SJ: Photoactivated gamma-secretase inhibitors directed to the active site covalently label presenilin 1. *Nature* 2000, 405:689–694
 13. Kornilova AY, Das C, Wolfe MS: Differential effects of inhibitors on the gamma-secretase complex. Mechanistic implications. *J Biol Chem* 2003, 278:16470–16473
 14. Hodgson G, Hager JH, Volik S, Hariono S, Wernick M, Moore D, Nowak N, Albertson DG, Pinkel D, Collins C, Hanahan D, Gray JW: Genome scanning with array CGH delineates regional alterations in mouse islet carcinomas. *Nat Genet* 2001, 29:459–464
 15. Tang YC, Amon A: Gene copy-number alterations: a cost-benefit analysis. *Cell* 2013, 152:394–405
 16. Huang X, Gollin SM, Raja S, Godfrey TE: High-resolution mapping of the 11q13 amplicon and identification of a gene, TAOS1, that is amplified and overexpressed in oral cancer cells. *Proc Natl Acad Sci U S A* 2002, 99:11369–11374
 17. Hyman E, Kauraniemi P, Hautaniemi S, Wolf M, Mousset S, Rozenblum E, Ringnér M, Sauter G, Monni O, Elkahoul A, Kallioniemi OP, Kallioniemi A: Impact of DNA amplification on gene expression patterns in breast cancer. *Cancer Res* 2002, 62:6240–6245
 18. Kusano N, Shiraishi K, Kubo K, Oga A, Okita K, Sasaki K: Genetic aberrations detected by comparative genomic hybridization in hepatocellular carcinomas: their relationship to clinicopathological features. *Hepatology* 1999, 29:1858–1862
 19. Midorikawa Y, Tsutsumi S, Nishimura K, Kamimura N, Kano M, Sakamoto H, Makuuchi M, Aburatani H: Distinct chromosomal bias of gene expression signatures in the progression of hepatocellular carcinoma. *Cancer Res* 2004, 64:7263–7270
 20. Andersson ER, Sandberg R, Lendahl U: Notch signaling: simplicity in design, versatility in function. *Development* 2011, 138:3593–3612
 21. Vooijs M, Liu Z, Kopan R: Notch: architect, landscaper, and guardian of the intestine. *Gastroenterology* 2011, 141:448–459
 22. Turnpenny PD, Ellard S: Alagille syndrome: pathogenesis, diagnosis and management. *Eur J Hum Genet* 2012, 20:251–257
 23. Sekiya S, Suzuki A: Intrahepatic cholangiocarcinoma can arise from Notch-mediated conversion of hepatocytes. *J Clin Invest* 2012, 122:3914–3918
 24. Benedito R, Roca C, Sörensen I, Adams S, Gossler A, Fruttiger M, Adams RH: The notch ligands Dll4 and Jagged1 have opposing effects on angiogenesis. *Cell* 2009, 137:1124–1135
 25. Doody RS, Raman R, Farlow M, Iwatsubo T, Vellas B, Joffe S, Kieburtz K, He F, Sun X, Thomas RG, Aisen PS; Alzheimer's Disease Cooperative Study Steering Committee, Siemers E, Sethuraman G, Mohs R; Semagacestat Study Group: A phase 3 trial of semagacestat for treatment of Alzheimer's disease. *N Engl J Med* 2013, 369:341–350
 26. He G, Luo W, Li P, Remmers C, Netzer WJ, Hendrick J, Bettayeb K, Flajolet M, Gorelick F, Wennogle LP, Greengard P: Gamma-secretase activating protein is a therapeutic target for Alzheimer's disease. *Nature* 2010, 467:95–98
 27. Maraver A, Fernandez-Marcos PJ, Herranz D, Cañamero M, Muñoz-Martin M, Gómez-López G, Mulero F, Megías D, Sanchez-Carbayo M, Shen J, Sanchez-Céspedes M, Palomero T, Ferrando A, Serrano M: Therapeutic effect of γ -secretase inhibition in KrasG12V-driven non-small cell lung carcinoma by derepression of DUSP1 and inhibition of ERK. *Cancer Cell* 2012, 22:222–234
 28. Lobry C, Ntziachristos P, Ndiaye-Lobry D, Oh P, Cimmino L, Zhu N, Araldi E, Hu W, Freund J, Abdel-Wahab O, Ibrahim S, Skokos D, Armstrong SA, Levine RL, Park CY, Aifantis I: Notch pathway activation targets AML-initiating cell homeostasis and differentiation. *J Exp Med* 2013, 210:301–319
 29. Kannan S, Sutphin RM, Hall MG, Golfman LS, Fang W, Nolo RM, Akers LJ, Hammit RA, McMurray JS, Kornblau SM, Melnick AM, Figueroa ME, Zweidler-McKay PA: Notch activation inhibits AML growth and survival: a potential therapeutic approach. *J Exp Med* 2013, 210:321–337
 30. López-Guerra M, Xargay-Torrent S, Rosich L, Montraveta A, Roldán J, Matas-Céspedes A, Villamor N, Aymerich M, López-Otín C, Pérez-Galán P, Roué G, Campo E, Colomer D: The γ -secretase inhibitor PF-03084014 combined with fludarabine antagonizes migration, invasion and angiogenesis in NOTCH1-mutated CLL cells. *Leukemia* 2015, 29:96–106
 31. Thorgeirsson SS, Lee JS, Grisham JW: Functional genomics of hepatocellular carcinoma. *Hepatology* 2006, 43:S145–S150
 32. Lee JS, Heo J, Libbrecht L, Chu IS, Kaposi-Novak P, Calvisi DF, Mikaelyan A, Roberts LR, Demetris AJ, Sun Z, Nevens F, Roskams T, Thorgeirsson SS: A novel prognostic subtype of human hepatocellular carcinoma derived from hepatic progenitor cells. *Nat Med* 2006, 12:410–416
 33. Wang F, Xia X, Wang J, Sun Q, Luo J, Cheng B: Notch1 signaling contributes to the oncogenic effect of HBx on human hepatic cells. *Biotechnol Lett* 2013, 35:29–37
 34. Tschaharganeh DF, Chen X, Latzko P, Malz M, Gaida MM, Felix K, Ladu S, Singer S, Pinna F, Gretz N, Sticht C, Tomasi ML, Delogu S, Evert M, Fan B, Ribback S, Jiang L, Brozzetti S, Bergmann F, Dombrowski F, Schirmacher P, Calvisi DF, Breuhahn K: Yes-associated protein up-regulates Jagged-1 and activates the Notch pathway in human hepatocellular carcinoma. *Gastroenterology* 2013, 144:1530–1542.e12
 35. Scaltriti M, Eichhorn PJ, Cortés J, Prudkin L, Aura C, Jiménez J, Chandralapaty S, Serra V, Prat A, Ibrahim YH, Guzmán M, Gili M, Rodríguez O, Rodríguez S, Pérez J, Green SR, Mai S, Rosen N, Hudis C, Baselga J: Cyclin E amplification/overexpression is a

- mechanism of trastuzumab resistance in HER2+ breast cancer patients. *Proc Natl Acad Sci U S A* 2011, 108:3761–3766
36. Wei W, Mok SC, Oliva E, Kim SH, Mohapatra G, Birrer MJ: FGF18 as a prognostic and therapeutic biomarker in ovarian cancer. *J Clin Invest* 2013, 123:4435–4448
 37. Ahn SM, Jang SJ, Shim JH, Kim D, Hong SM, Sung CO, Baek D, Haq F, Ansari AA, Lee SY, Chun SM, Choi S, Choi HJ, Kim J, Kim S, Hwang S, Lee YJ, Lee JE, Jung WR, Jang HY, Yang E, Sung WK, Lee NP, Mao M, Lee C, Zucman-Rossi J, Yu E, Lee HC, Kong G: A genomic portrait of resectable hepatocellular carcinomas: implications of RB1 and FGF19 aberrations for patient stratification. *Hepatology* 2014, 60:1972–1982
 38. Wang K, Lim HY, Shi S, Lee J, Deng S, Xie T, Zhu Z, Wang Y, Pocalyko D, Yang WJ, Rejto PA, Mao M, Park CK, Xu J: Genomic landscape of copy number aberrations enables the identification of oncogenic drivers in hepatocellular carcinoma. *Hepatology* 2013, 58: 706–717
 39. Hurwitz H, Fehrenbacher L, Novotny W, Cartwright T, Hainsworth J, Heim W, Berlin J, Baron A, Griffing S, Holmgren E, Ferrara N, Fyfe G, Rogers B, Ross R, Kabbinavar F: Bevacizumab plus irinotecan, fluorouracil, and leucovorin for metastatic colorectal cancer. *N Engl J Med* 2004, 350:2335–2342
 40. Hottinger AF, Aissa AB, Espeli V, Squiban D, Dunkel N, Vargas MI, Hundsberger T, Mach N, Schaller K, Weber DC, Bodmer A, Dietrich PY: Phase I study of sorafenib combined with radiation therapy and temozolomide as first-line treatment of high-grade glioma. *Br J Cancer* 2014, 110:2655–2661
 41. Bachelot T, Garcia-Saenz JA, Verma S, Gutierrez M, Pivot X, Kozloff MF, Prady C, Huang X, Khosravan R, Wang Z, Cesari R, Tassell V, Kern KA, Blay JY, Lluch A: Sunitinib in combination with trastuzumab for the treatment of advanced breast cancer: activity and safety results from a phase II study. *BMC Cancer* 2014, 14:166
 42. Yamashita T, Forgues M, Wang W, Kim JW, Ye Q, Jia H, Budhu A, Zanetti KA, Chen Y, Qin LX, Tang ZY, Wang XW: EpCAM and alpha-fetoprotein expression defines novel prognostic subtypes of hepatocellular carcinoma. *Cancer Res* 2008, 68:1451–1461
 43. Noel PR, Barnett KC, Davies RE, Jolly DW, Leahy JS, Mawdesley-Thomas LE, Shillam KW, Squires PF, Street AE, Tucker WC, Worden AN: The toxicity of dimethyl sulphoxide (DMSO) for the dog, pig, rat and rabbit. *Toxicology* 1975, 3:143–169