

Commentary

Mechanogenomic Control of DNA Exposure and Sequestration

Gary S. Stein

From the University of Massachusetts Medical School,
Worcester, Massachusetts

There is longstanding recognition and appreciation for striking differences in the structure, assembly and organization of regulatory machinery in normal and cancer cells. Drawings from 100 years ago of Galeotti's comparisons of cancer cell and normal cell chromosomes illustrated gross differences in chromatin structure and chromosome numbers in human tissue harboring a tumor.¹ Tumor-related perturbations in nuclear structure-function interrelationships are well documented by prominent perturbations in the size and number of nucleoli, indicating the reconfiguration of the nuclear infrastructure that supports ribosomal gene expression and consequently protein biosynthesis. However, how higher order chromatin structure is controlled and maintained by the cancer cell and its microenvironment, compared to normal cells, has remained beyond our reach. The work presented by Maniotis et al² in this issue of *The American Journal of Pathology* shows there are striking differences in restriction enzyme sensitivity between tumor cell and normal cell genomes. Furthermore, Maniotis et al propose that the mechanical contiguity and coordination of nuclear proteins bound to DNA, the cytoskeleton, and the extracellular matrix (ECM) may work in concert to provide a cytoarchitectural resistance mechanism deep within the cell. When verified in other laboratories, these findings may constitute the basis for a paradigm shift in the way we view how the genomes of higher eukaryotic cells, and malignant cells in particular, are regulated.

By using DNase digestions and nick-end labeling techniques, a previous generation of tumor biologists had established that in order for a gene to become expressed, it had to be "exposed." Moreover, Puck et al³ had shown that transformed and tumor cells that were "reverse transformed" to a normal phenotype with various chemical compounds exhibited a shift in their nuclei's sensitivity to DNase I, and exhibited profound changes in their cytoskeletons and overall morphology.

Maniotis et al² have compared the extent of exposure or sequestration of a well-characterized collection of highly invasive, poorly invasive, and normal human cell genomes to digestion by specific restriction enzymes.

Restriction enzyme digests performed on intact nuclei, not treated with any other chemical agents, showed that *AluI* and *MspI* both distinguished normal cell genomes from tumor cell genomes. For example, highly invasive cell nuclei always resisted digestion with *AluI* and *MspI* compared to poorly invasive or normal cells in permeabilized cell models, in cell smear preparations, in suspended cells using flow cytometry, and in touch preps of human tissue. Specificity is suggested because other restriction enzymes did not discriminate among normal or tumor cell genomes.

Differences in Alu site exposure were shown to be independent of the cell cycle, as shown by comparing digestions of complete sets of mitotic chromosomes^{4,5} with digestions of interphase nuclei from the same cell types. Taken together with other restriction sites that exhibit cell cycle-stage specific accessibility⁶⁻⁸ it appears that there are both conserved and transient components of chromatin organization that must be accommodated during the cell cycle period. It is necessary to mechanistically explain parameters of mitotic chromosome condensation that are independent from the sequestration of restriction sites along the DNA, and the observation that nuclear matrix-associated AML⁹⁻¹¹ and ALL^{12,13} phenotype-restricted transcription factors, as well as UBF-1¹⁴ remain on target gene loci during mitosis. New models are required to explain higher order chromatin structure more comprehensively. Such models should account for components of cell cycle-driven chromatin condensation that are operative independent of how specific restriction enzyme sites are cryptic or exposed in highly invasive, poorly invasive, and normal cells.

Experiments were also presented in the study that showed exposure of chromatin or chromosomes to DTT or β -mercaptoethanol rendered highly invasive and sequestered cell genomes sensitive to *AluI*. This suggests that proteins rich in disulfide bonds are involved in the sequestration phenomenon in highly invasive cell nuclei.

Accepted for publication January 21, 2005.

This commentary relates to Maniotis et al, *Am J Pathol* 2005, 166:1187-1203, published in this issue.

Address reprint requests to Gary Stein, Department of Cell Biology, University of Massachusetts Medical School, 55 Lake Ave. North, Worcester, MA 01655. E-mail: gary.stein@umassmed.edu.

Enzymes or reagents that remove histones or topoisomerase⁵ did not discriminate among genomes tested from different cell types. These observations are consistent with the idea that alterations in higher order chromatin structure and may, in part, be controlled by disulfide-rich nuclear matrix proteins,¹⁵⁻²¹ which orchestrate widespread changes in restriction enzyme site exposure and sequestration. However, these digestion experiments also suggest that the disulfide-rich nuclear matrix cannot be a static or fixed structure, which is consistent with current models of chromatin structure and higher order nuclear organization of gene regulatory machinery in nuclear matrix-associated microenvironments that support combinatorial control of gene expression by dynamic, physiologically responsive targeting and retention mechanisms.²²⁻³⁴

Because digestions of DNA in nuclei of normal cells with *AluI* and *MspI* appear to additionally separate nucleoli from structural components of the nucleus, it appears that the DNA within normal cell chromatin, when it is not digested, may impose constraints that contribute to the architectural organization of intranuclear structure. Maniotis' case² for DNA as a principal scaffolding on which nuclear proteins are bound, and not vice versa, is consistent with his previous observations of nuclease digestions of micro surgically isolated mitotic chromosomes derived from normal endothelial cells and fibroblasts,⁴ and is supported by his observation that complete chromosome sets are produced as intact genomes from both normal mitotic cells and mitotic tumor cells via microsurgical removal of any single chromosome.^{4,5} Reversal of the relative insensitivity of chromatin of highly invasive cells to *AluI* by disulfide-disrupting agents suggests that the sensitivity of invasive nuclei to *AluI* and disulfide-rich protein disruption may be due to increased representation of these disulfide-rich proteins, or may suggest that there is a higher affinity of these proteins for the DNA of highly invasive cells. Further work is needed to determine whether nicking the DNA with *AluI* leads to discontinuities along the DNA, which in turn permits DNA supercoiling to unravel sufficiently to dissociate the disulfide-rich proteins from the DNA throughout the genome, and allow complete digestion to occur.

There is a requirement to characterize the nuclear matrix disulfide-rich proteins that may be facilitating architecturally linked control of gene expression. Equally important is the necessity to relate the involvement of DNA as a scaffold for control of gene expression that differs in normal and cancer cells with the growing evidence for autologous intranuclear trafficking of regulatory proteins^{11,32-34} that are key for organization and assembly of the regulatory machinery for context-dependent, combinatorial control of gene expression in nuclear matrix-associated microenvironments of interphase cells and at chromosomal loci of mitotic cells.³⁵⁻⁴³

In the second part of the work presented by Maniotis et al² in this issue, a systematic search was initiated to test if growth factors, or soluble and polymerized ECM molecules, affect the sequestration or exposure of DNA, and to test if there was specificity of different ECM molecule types on Alu site exposure or sequestration. Exposure of

living cells to soluble laminin and RGD-C resulted in sequestration of Alu sites from digestion by *AluI* in all cells tested. However, sequestration induced by Matrigel, laminin, and RGD-C was always more intense in the nuclei of cells of increasing grades of invasiveness. Serum, fibronectin, bFGF, EGF, and type I collagen had no effect on the sequestration of *AluI* sensitive sites. When cells were grown in or on polymerized Matrigel or laminin platforms, profound sequestration was observed. This was in contrast to cells that were situated on Type I collagen, serum, or fibronectin. In addition, when comparing six repeated microarrays of highly invasive cells situated on Matrigel versus the same highly invasive cells not situated on Matrigel, a consistent differential expression of 990 transcripts was obtained. These results, when taken together with the observed chemical changes in sensitivity underlying the nuclease digestions, provide preliminary evidence that exposure of only one extracellular matrix molecule on the cell surface for 30 minutes can induce global effects on chromatin organization.

In multicellular organisms, the cytoskeleton and extracellular matrix are known to play a fundamental role in determining cellular behaviors. To test if different cytoskeletal fiber systems influenced the sequestration or exposed the Alu sites, a variety of cytoskeleton-disrupting drugs were used by Maniotis et al² to determine whether the higher order structure of chromatin was controlled by actin, microtubules, or intermediate filaments. Maniotis et al⁴⁴ had previously shown that a tug on an integrin receptor could alter the molecular alignment of intranuclear molecules in 1 second, and that each cytoskeletal system exerted a different stabilizing effect on both nuclear structure, and force transduction. In this study, each cytoskeletal fiber system profoundly contributed to the sequestration or exposure effect. Actin disruption decreased sequestration, while microtubule or intermediate filament disruption dramatically increased sequestration. In these experiments, the data suggest that nuclear size and cytoplasmic spreading and gene sequestration or exposure are co-events. A similar relationship between cell growth, cell "differentiation," and nuclear size in endothelial cells was shown long ago by Folkman and Moscona⁴⁵ on monolayers coated with increasing amounts of a cellular adhesive. The results of the Maniotis et al study² extend these observations to include the sequestration and exposure of specific DNA sites, and a compelling and testable model is advanced that involves cytoskeletal control of nuclear structure and nuclear pore complexes.

Together, these findings raise, and experimentally address, new questions regarding the fundamental way genetic information is controlled by proteins containing disulfide-rich bonds, by the extracellular matrix microenvironment, and by the cytoskeleton. The work also has contributed several new approaches that can be used as diagnostic tools. Furthermore, the observed differences in the sequestration of DNA among cells of varying invasive behavior can be exploited to distinguish differing degrees of malignancy that do not depend on presently used molecular markers. Therefore, the differential sensitivity of Alu sites in highly invasive tumor cells versus

noninvasive tumor cells offers an alternative for dependence on molecular markers, which have been shown to be highly variable in the context of vasculogenic mimicry, and which manifests as molecular mimicry (deregulated protein expression) in tissue sections of the most invasive types of tumors.^{46–48} For example, the most highly invasive melanoma tumor cells, although they ultimately derive from the neural crest, are able to express protein markers characteristic of a variety of different normal cell types (such as endothelial cells), but also markers specific to epithelial cells, and a wide variety of other cell types, depending on the degree of deregulation. In this context, clinically relevant lessons can be learned about mechanisms for drug resistance that are mediated by microenvironments that are key components of tissue organization.^{49,50}

All considered, it appears that how genes are sequestered, exposed, and expressed in higher eukaryotic cells can be viewed as part of a mechanically coordinated, cell shape-dependent, hierarchical system.^{49,51–61} In this regard, it has been proposed that the principles of tensional integrity (tensegrity) may be a useful conceptual framework to explore and predict how molecules can function collectively as components of integrated, hierarchical systems, in the physical context of normal living cells, and tissues.⁶²

If the information directing these processes are indeed “mechanogenomic” in nature, as described by Maniotis et al² in this issue of *The American Journal of Pathology*, the apparent complexity of development and cancer may be better understood by insight into how the genome is part of a larger machine that can interpret different signals only in a limited number of ways that is restricted by the ECM microenvironment, and the type of mechanogenomic architecture that environment erects within the cell and nucleus.

In a broader context, understanding of cell structure-gene expression interrelationships is in its infancy. However, there is accruing evidence for a central role of cellular microenvironments in the detection, integration, and execution of regulatory signals. Support for focal thresholds of genes, transcripts, receptors, and regulatory factors that optimize informative encounters to support gene expression, replication, and repair at multiple levels has emerged from interrogation of fundamental parameters of biological control. Additional insight into the regulation of gene expression should be forthcoming from further exploration of the mechanisms and underlying parameters that mediate the multistep dynamic positioning and combinatorial association of architecturally organized regulatory macromolecules in the nucleus, cytoplasm, and extracellular matrix. Despite compelling support for the physiological relevance of intracellular regulatory domains, there is a requirement to define rate limiting parameters of mechanisms that mediate the temporal and spatial components of cell structure-gene expression interrelationships. The rules governing organization of the regulatory machinery in the three-dimensional context of cellular architecture are being functionally enhanced by the combined applications of molecular, cellular, and *in vivo* genetic approaches. A prominent position for mechanogenomic control is an

important consideration. It is realistic to expect that mechanistic explanations for the dynamic organization, assembly, and activities of regulatory machinery in cellular microenvironments which are subtly or catastrophically compromised in many diseases, including cancer, can provide a platform for novel approaches to diagnosis and treatment. It is further necessary to incorporate consideration of mechanogenomic control into mechanisms that may be operative in physiological processes that include, but are not restricted to, cell motility that is coupled with development, differentiation, tissue turnover, and remodeling, as well as the common denominators that may link mechanogenomic components of control with tumor progression and metastasis.

References

1. Wilson EB: Pathological mitosis in cancer cells. *The Cell in Development and Inheritance. The Sources of Science No. 30.* New York and London, Johnson Reprint Corporation, 1966, pp 68–69
2. Maniotis AJ, Valyi-Nagy K, Karavitis J, Moses J, Boddipali V, Wang Y, Nuñez R, Setty S, Arbieva Z, Bissell MJ, Folberg R: Chromatin organization measured by Alu I restriction enzyme changes with malignancy and is regulated by the extracellular matrix and the cytoskeleton. *Am J Pathol* 2005, 166:1187–1203
3. Puck TT, Krystosek A, and Chan DC: Genome regulation in mammalian cells. *Somat Cell Mol Genet* 1990, 16:257–265
4. Maniotis AJ, Bojanowski K, Ingber DE: Mechanical continuity and reversible chromosome disassembly within intact genomes removed from living cells. *J Cell Biochem* 1997, 65:114–130
5. Bojanowski K, Maniotis AJ, Plisov S, Larsen AK, Ingber DE: DNA topoisomerase II can drive changes in higher order chromosome architecture without enzymatically modifying DNA. *J Cell Biochem* 1998, 69:127–142
6. Chrysogelos S, Riley DE, Stein G, Stein J: A human histone H4 gene exhibits cell cycle-dependent changes in chromatin structure that correlate with its expression. *Proc Natl Acad Sci USA* 1985, 82:7535–7539
7. Chrysogelos S, Pauli U, Stein G, Stein J: Fine mapping of the chromatin structure of a cell cycle-regulated human H4 histone gene. *J Biol Chem* 1989, 264:1232–1237
8. Stein G, Park W, Thrall C, Mans R, Stein J: Regulation of cell cycle stage-specific transcription of histone genes from chromatin by non-histone chromosomal proteins. *Nature* 1975, 257:764–767
9. Zaidi SK, Young DW, Pockwinse SH, Javed A, Lian JB, Stein JL, van Wijnen AJ, Stein GS: Mitotic partitioning and selective reorganization of tissue specific transcription factors in progeny cells. *Proc Natl Acad Sci USA* 2003, 100:14852–14857
10. Young DW, Zaidi SK, Furcinitti PS, Javed A, van Wijnen AJ, Stein JL, Lian JB, Stein GS: Quantitative signature for architectural organization of regulatory factors using intranuclear informatics. *J Cell Sci* 2004, 117:4889–4896
11. Zeng C, van Wijnen AJ, Stein JL, Meyers S, Sun W, Shopland L, Lawrence JB, Penman S, Lian JB, Stein GS, Hiebert SW: Identification of a nuclear matrix targeting signal in the leukemia and bone-related AML/CBF α transcription factors. *Proc Natl Acad Sci USA* 1997, 94:6746–6751
12. Ennas MG, Sorio C, Greim R, Nieddu M, Scarpa A, Orlandini S, Croce CM, Fey GH, Marschalek R: The human ALL-1/MLL/HRX antigen is predominantly localized in the nucleus of resting and proliferating peripheral blood mononuclear cells. *Cancer Res* 1997, 57:2035–2041
13. Nakamura T, Mori T, Tada S, Krajewski W, Rozovskaia T, Wassell R, Dubois G, Mazo A, Croce CM, Canaani E: ALL-1 is a histone methyltransferase that assembles a supercomplex of proteins involved in transcriptional regulation. *Mol Cell* 2002, 10:1119–1128
14. Gebrane-Younes J, Fomproix N, Hernandez-Verdun D: When rDNA transcription is arrested during mitosis, UBF is still associated with non-condensed rDNA. *J Cell Sci* 1997, 110:2429–2440

15. Berezney R, Coffey DS: Nuclear protein matrix: association with newly synthesized DNA. *Science* 1975, 189:291–292
16. Capco DG, Wan KM, Penman S: The nuclear matrix: three-dimensional architecture and protein composition. *Cell* 1982, 29:847–858
17. Pienta KJ, Coffey DS: A structural analysis of the role of the nuclear matrix and DNA loops in the organization of the nucleus and chromosome. *J Cell Sci Suppl* 1984, 1:123–135
18. Nickerson JA, Krockmalnic G, Wan KM, Turner CD, Penman S: A normally masked nuclear matrix antigen that appears at mitosis on cytoskeleton filaments adjoining chromosomes, centrioles and midbodies. *J Cell Biol* 1992, 116:977–987
19. Mancini MA, He D, Ouspenski II, Brinkley BR: Dynamic continuity of nuclear and mitotic matrix proteins in the cell cycle. *J Cell Biochem* 1996, 62:158–164
20. Roti Roti JL, Wright WD, VanderWaal R: The nuclear matrix: a target for heat shock effects and a determinant for stress response. *Crit Rev Eukaryot Gene Expr* 1997, 7:343–360
21. Stein GS, Roberts RM, Davis JL, Head WJ, Stein JL, Thrall CL, Van Veen J, Welch DW: Are glycoproteins and glycosaminoglycans components of the eukaryotic genome? *Nature* 1975, 258:639–641
22. Stein GS, Montecino M, van Wijnen AJ, Stein JL, Lian JB: Nuclear structure - gene expression interrelationships: implications for aberrant gene expression in cancer. *Cancer Res* 2000, 60:2067–2076
23. Zaidi SK, Young DW, Choi JY, Pratap J, Javed A, Montecino M, Stein JL, Lian JB, van Wijnen AJ, Stein GS: Intracellular trafficking: organization and assembly of regulatory machinery for combinatorial biological control. *J Biol Chem* 2004, 279:43363–43366
24. Stein GS, Zaidi SK, Braastad CD, Montecino M, van Wijnen AJ, Choi J-Y, Stein JL, Lian JB, Javed A: Functional architecture of the nucleus: organizing the regulatory machinery for gene expression, replication and repair. *Trends Cell Biol* 2003, 13:584–592
25. Zink D, Fischer AH, Nickerson JA: Nuclear structure in cancer cells. *Nat Rev Cancer* 2004, 4:677–687
26. Misteli T: The concept of self-organization in cellular architecture. *J Cell Biol* 2001, 155:181–185
27. Olson MO, Hingorani K, Szebeni A: Conventional and nonconventional roles of the nucleolus. *Internat Rev Cytol* 2002, 219:199–266
28. Spencer VA, Davie JR: Signal transduction pathways and chromatin structure in cancer cells. *J Cell Biochem Suppl* 2000, Suppl 35:27–35
29. Stachowiak MK, Fang X, Myers JM, Dunham SM, Berezney R, Maher PA, Stachowiak EK: Integrative nuclear FGFR1 signaling (INFS) as a part of a universal “feed-forward-and-gate” signaling module that controls cell growth and differentiation. *J Cell Biochem* 2003, 90:662–691
30. Taatjes DJ, Marr MT, Tjian R: Regulatory diversity among metazoan co-activator complexes. *Nat Rev Mol Cell Biol* 2004, 5:403–410
31. Kosak ST, Groudine M: Gene order and dynamic domains. *Science* 2004, 306:644–647
32. DeFranco DB: Navigating steroid hormone receptors through the nuclear compartment. *Mol Endocrinol* 2002, 16:1449–1455
33. Nagaich AK, Rayasam GV, Martinez ED, Becker M, Qiu Y, Johnson TA, Elbi C, Fletcher TM, John S, Hager GL: Subnuclear trafficking and gene targeting by steroid receptors. *Ann NY Acad Sci* 2004, 1024:213–220
34. Phair RD, Scaffidi P, Elbi C, Vecerova J, Dey A, Ozato K, Brown DT, Hager G, Bustin M, Misteli T: Global nature of dynamic protein-chromatin interactions in vivo: three-dimensional genome scanning and dynamic interaction networks of chromatin proteins. *Mol Cell Biol* 2004, 24:6393–6402
35. Cook PR: The organization of replication and transcription. *Science* 1999, 284:1790–1795
36. Choi J-Y, Pratap J, Javed A, Zaidi SK, Xing L, Balint E, Dalamangas S, Boyce B, van Wijnen AJ, Lian JB, Stein JL, Jones SN, Stein GS: Subnuclear targeting of Runx/Cbfa/AML factors is essential for tissue-specific differentiation during embryonic development. *Proc Natl Acad Sci USA* 2001, 98:8650–8655
37. Brown KE, Baxter J, Graf D, Merkschlager M, Fisher AG: Dynamic repositioning of genes in the nucleus of lymphocytes preparing for cell division. *Mol Cell* 1999, 3:207–217
38. Ma H, Siegel AJ, Berezney R: Association of chromosome territories with the nuclear matrix: disruption of human chromosome territories correlates with the release of a subset of nuclear matrix proteins. *J Cell Biol* 1999, 146:531–542
39. Javed A, Barnes GL, Pratap J, Antkowiak T, Gerstenfeld LC, van Wijnen AJ, Stein JL, Lian JB, Stein GS: Impaired intranuclear trafficking of Runx2 (AML3/CBFA1) transcription factors in breast cancer cells inhibits formation of osteolytic lesions in vivo. *Proc Natl Acad Sci USA* 2005, 102:1454–1459
40. Misteli T, Spector DL: RNA polymerase II targets pre-mRNA splicing factors to transcription sites in vivo. *Mol Cell* 1999, 3:697–705
41. Scully R, Livingston DM: In search of the tumour-suppressor functions of BRCA1 and BRCA2. *Nature* 2000, 408:429–432
42. Smith KP, Moen PT, Wydner KL, Coleman JR, Lawrence JB: Processing of endogenous pre-mRNAs in association with SC-35 domains is gene specific. *J Cell Biol* 1999, 144:617–629
43. Wagner S, Chiosea S, Nickerson JA: The spatial targeting and nuclear matrix binding domains of SRm160. *Proc Natl Acad Sci USA* 2003, 100:3269–3274
44. Maniatis AJ, Chen CS, Ingber DE: Demonstration of mechanical connections between integrins, cytoskeletal filaments, and nucleoplasm that stabilize nuclear structure. *Proc Natl Acad Sci USA* 1997, 94:849–854
45. Folkman J, Moscona A: Role of cell shape in growth control. *Nature* 1978, 273:345–349
46. Maniatis AJ, Folberg R, Hess A, Sefter EA, Gardner LM, Pe'er J, Trent JM, Meltzer PS, Hendrix MJ: Vascular channel formation by human melanoma cells in vivo and in vitro: vasculogenic mimicry. *Am J Pathol* 1999, 155:739–752
47. Chen X, Maniatis AJ, Majumdar D, Pe'er J, Folberg R: Uveal melanoma cell staining for CD34 and assessment of tumor vascularity. *Invest Ophthalmol Vis Sci* 2002, 43:2533–2539
48. Sefter EA, Meltzer PS, Kirschmann DA, Pe'er J, Maniatis AJ, Trent JM, Folberg R, Hendrix MJ: Molecular determinants of human uveal melanoma invasion and metastasis. *Clin Exp Metastasis* 2002, 19:233–246
49. Bissell MJ, Rizki A, Mian IS: Tissue architecture: the ultimate regulator of breast epithelial function. *Curr Opin Cell Biol* 2003, 15:753–762
50. Boudreau N, Myers C, Bissell MJ: From laminin to lamin: regulation of tissue-specific gene expression by the ECM. *Trends Cell Biol* 1995, 5:1–4
51. Itoh M, Bissell MJ: The organization of tight junctions in epithelia: implications for mammary gland biology and breast tumorigenesis. *J Mammary Gland Biol Neoplasia* 2003, 8:449–462
52. Novaro V, Radisky DC, Ramos Castro NE, Weisz A, Bissell MJ: Malignant mammary cells acquire independence from extracellular context for regulation of estrogen receptor alpha. *Clin Cancer Res* 2004, 10:402S–409S
53. Schmeichel KL, Bissell MJ: Modeling tissue-specific signaling and organ function in three dimensions. *J Cell Sci* 2003, 116:2377–2388
54. Roskelley CD, Srebrow A, Bissell MJ: A hierarchy of ECM-mediated signalling regulates tissue-specific gene expression. *Curr Opin Cell Biol* 1995, 7:736–747
55. Bissell MJ, Barcellos-Hoff MH: The influence of extracellular matrix on gene expression: is structure the message? *J Cell Sci Suppl* 1987, 8:327–343
56. Bissell MJ, Hall HG, Parry G: How does the extracellular matrix direct gene expression? *J Theor Biol* 1982, 99:31–68
57. Sood AK, Coffin JE, Schneider GB, Fletcher MS, DeYoung BR, Guman LM, Gershenson DM, Schaller MD, Hendrix MJ: Biological significance of focal adhesion kinase in ovarian cancer: role in migration and invasion. *Am J Pathol* 2004, 165:1087–1095
58. Chunthapong J, Sefter EA, Khalkhali-Ellis Z, Sefter RE, Amir S, Lubaroff DM, Heidger PM, Jr., Hendrix MJ: Dual roles of E-cadherin in prostate cancer invasion. *J Cell Biochem* 2004, 91:649–661
59. Sood AK, Hendrix MJ: The complexity of tumor vascularity. *Cancer Biol Ther* 2003, 2:257–258
60. Hendrix MJ, Sefter EA, Kirschmann DA, Quaranta V, Sefter RE: Remodeling of the microenvironment by aggressive melanoma tumor cells. *Ann NY Acad Sci* 2003, 995:151–161
61. Hendrix MJ, Sefter EA, Kirschmann DA, Sefter RE: Molecular biology of breast cancer metastasis: molecular expression of vascular markers by aggressive breast cancer cells. *Breast Cancer Res* 2000, 2:417–422
62. Ingber DE: Tensegrity II: how structural networks influence cellular information processing networks. *J Cell Sci* 2003, 116:1397–1408